arXiv:1109.1233v1 [math.PR] 6 Sep 2011

Cycle structure of percolation on high-dimensional tori

Remco van der Hofstad [*] Artëm Sapozhnikov [†]

September 7, 2011

Abstract

In the past years, many properties of the critical behavior of the largest connected components on the high-dimensional torus, such as their sizes and diameter, have been established. The order of magnitude of these quantities equals the one for percolation on the complete graph or Erdős-Rényi random graph, raising the question whether the scaling limits or the largest connected components, as identified by Aldous (1997), are also equal.

In this paper, we investigate the *cycle structure* of the largest critical components for high-dimensional percolation on the torus $\{1, \dots, r\}^d$. While percolation clusters naturally have many *short* cycles, we show that the *long* cycles, i.e., cycles that pass through the boundary of the cube of width $r/2$ centered around each of their vertices, have length of order $r^{d/3}$, as on the critical Erdős-Rényi random graph. On the Erdős-Rényi random graph, cycles play an essential role in the scaling limit of the large critical clusters, as identified by Addario-Berry, Broutin and Goldschmidt (2010).

Our proofs crucially rely on extensions of results of Kozma and Nachmias (2011) about the one-arm exponent for critical high-dimensional percolation. We show that, at criticality, the length of an open path from the origin to (Euclidean) distance n, if it exists, is of order n^2, which is the same as for critical branching random walk.

Key words: Random graph, phase transition, critical behavior, percolation, cycle structure.
MSC2000 subject classification. 05C80, 60K35, 82B43.

1 Introduction and results

In the past years, the investigation of percolation on various high-dimensional tori has attracted tremendous attention. In [5, 6], the phase transition of the largest connected component was investigated for percolation on general high-dimensional tori, including the complete graph, the hyper-cube in high dimensions, as well as finite-range percolation in sufficiently high dimensions. The phase transition of percolation on high-dimensional tori is *mean-field*, i.e., it shares many features with that on the complete graph as identified in [9] (see, e.g., [2, 4, 16, 17, 21]).

In [5], the subcritical and critical behavior was investigated under the so-called *triangle condition*, a general assumption on the underlying graph that ensures that the model is mean-field. The critical behavior of the model was identified in terms of the blow-up of the expected cluster size, which identifies a window of critical values of the edge occupation probabilities. For any parameter value in this critical

---

[*] Department of Mathematics and Computer Science, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands. E-mail: `rhofstad@win.tue.nl`

[†] ETH Zürich, Department of Mathematics, Rämistrasse 101, 8092 Zürich. E-mail: `artem.sapozhnikov@math.ethz.ch`

window, the largest connected component was shown to be of order $V^{2/3}$, as on the complete graph, where V denotes the number of vertices in the graph. In [6], the triangle condition was proved to hold for the above-mentioned examples.

The situation of finite-range high-dimensional tori, which in the graph sense converge to the hypercubic lattice, was brought substantially further in [14, 15], where, among others, it was shown that the percolation critical value on the infinite lattice lies inside the scaling window. We now know that the largest connected components are all of order $V^{2/3}$, that the maximal connected component $|\mathcal{C}_{\max}|$ satisfies that $|\mathcal{C}_{\max}|V^{-2/3}$ and $V^{2/3}/|\mathcal{C}_{\max}|$ are tight sequences of random variables that are non-concentrated, and that the diameter of large clusters is of order $V^{1/3}$. These results (and more) are also known to hold on the Erdős-Rényi random graph, see e.g., [2, 22], as well as the monographs [4, 17]. This raises the question whether the scaling limits agree.

1.1 Percolation in high dimensions

We consider bond percolation on a graph G. For a given parameter $p \in [0,1]$, this is the probability measure $\mathbb{P}_p$ on subgraphs of G defined as follows. We delete edges of G with probability $(1-p)$ and otherwise keep them, independently for different edges. The edges of the resulting random subgraph of G are called *open* and the deleted edges are called *closed*. Connected components of this random subgraph are called *open clusters*. The graphs we investigate in this paper are (a) the d-dimensional torus $\mathbb{T}_r^d = \{0,\dots,r-1\}^d$; and (b) the hypercubic lattice $\mathbb{Z}^d$, where the dimension d is supposed to be sufficiently large. How large we need to take d depends on the edge structure of G. We consider two different settings: (a) In the *nearest-neighbor model*, two vertices are connected by an edge if they are nearest-neighbors on G. With our choice of G, every vertex has $2d$ nearest-neighbors. In this setting we take the dimension d large enough. (b) In the *spread-out model* with a parameter L, two vertices are connected by an edge if there is a hypercube of size L in G that contains these vertices. With our choice of G, every vertex has $(2L+1)^d - 1$ neighbors. Of course, we are only interested in the case when the size of the torus is much larger than L. In the spread-out setting with large enough L, we take the dimension $d > 6$.

To justify our choice of dimension, we recall a number of well-known results about percolation on $\mathbb{Z}^d$. For bond percolation on $\mathbb{Z}^d$ with $d > 1$, there exists a critical probability $p_c \in (0,1)$ such that, for $p < p_c$, all open clusters are almost surely finite and, for $p > p_c$, there is almost surely an infinite open cluster. At $p = p_c$, it is widely believed that there is almost surely no infinite open cluster. This fact has been shown for $d = 2$ by Kesten [18] and for sufficiently large d by Barsky and Aizenman [3] and Hara and Slade [13]. Here, by sufficiently large d, we mean $d > 18$ for the nearest-neighbor model and $d > 6$ for the spread-out model with sufficiently large L. Showing this for all $d > 1$ still remains a challenging open problem.

The main assumption that we use in the paper concerns an estimate on the probability that, at criticality, two vertices x and y are in the same open cluster of bond percolation on $\mathbb{Z}^d$, which we denote by $x \leftrightarrow y$. We assume that there exist constants D_1 and D_2 such that, for all x and y in $\mathbb{Z}^d$,

$$D_1(1+|x-y|)^{2-d} \le \mathbb{P}_{p_c}(x \leftrightarrow y) \le D_2(1+|x-y|)^{2-d}. \tag{1.1}$$

These bounds have been established using so-called *lace expansion* techniques, for the nearest-neighbor model with large enough d by Hara [11], and for the spread-out model with $d > 6$ by Hara, van der Hofstad and Slade [12]. In fact, these papers give asymptotic formulas for such probabilities, but for our purposes, the bounds (1.1) suffice.

It is believed that the estimates (1.1) hold for the nearest-neighbor model with $d > 6$, however the proof of this fact is beyond the current methods. It has been proved by Chayes and Chayes [8] (assuming the existence of critical exponents) that the bounds (1.1) are violated for $d < 6$. The dimension $d = d_c = 6$ is usually referred to as the *upper critical dimension*.

A simple computation using the upper bound in (1.1) shows that

$$\nabla(p_c) = \sum_{x,y} \mathbb{P}_{p_c}(0 \leftrightarrow x)\mathbb{P}_{p_c}(x \leftrightarrow y)\mathbb{P}_{p_c}(y \leftrightarrow 0) < \infty. \tag{1.2}$$

The bound in (1.2) is called the *triangle condition*, and is believed to be true for $d > 6$. The triangle condition implies that the sub- and critical phases of percolation on $\mathbb{Z}^d$ behave similarly to the ones on a tree, for example, many critical exponents on $\mathbb{Z}^d$ are equal to those on the tree. Intuitively, the geometry of large critical clusters trivializes, since the space is so vast that far away clusters are close to being independent. In recent years, a related condition has been proved to hold on the torus, which implies that the critical behavior of large connected components on the high-dimensional torus is similar to that on the complete graph. Sometimes this is called *random graph asymptotics* for percolation on the high-dimensional torus.

In this paper, we study the cycle structure of bond percolation on the d-dimensional torus in the above two settings. Despite the fact that, for any $p \in (0, 1)$, the number of vertices in open cycles is dense in the torus, which is not the case for the critical and subcritical Erdős-Rényi random graph (see [21]), most of such vertices belong only to *short* cycles, such as open squares of four bonds. Short cycles vanish in the scaling limit of large critical clusters, and are thus irrelevant to the scaling limit. Therefore, we focus on the existence of open long cycles, where we say that a cycle is long when it passes through the boundary of the cube of width $r/2$ centered around each of its vertices. Special cases of long cycles are non-contractible cycles, which are cycles that cannot, when considered as continuous curves, be contracted to a point, and thus wind around the torus at least once.

Our main results show that the mean number of vertices in open long cycles grows like $V^{1/3}$, and that such cycles (when they exist) contain the order of $V^{1/3}$ vertices. Moreover, we show that the probability of the existence of at least one open long cycle in a large cluster is bounded away from 0 and 1, uniformly in the volume of the graph. As we discuss in more details below, this situation is analogous to the situation on the complete graph, as investigated in [1, 2, 21]. We also refer the reader to [21, pages 722-723] for the discussion of more refined results about the structure of connected components of the critical Erdős-Rényi random graph.

For simplicity of presentation, *we restrict ourselves hereafter to the nearest-neighbor model.* The results of this paper still hold for the spread-out model on the d-dimensional torus with $d > 6$.

The remainder of this section is organized as follows. In Section 1.2, we describe our main results, in Section 1.3 we describe some results on critical percolation on $\mathbb{Z}^d$ and the torus that are used in the proofs of our main results and are interesting in their own right, and in Section 1.4, we discuss the results and their relation to the work on the Erdős-Rényi random graph.

1.2 Main results

We start by introducing some notation. For $a \in \mathbb{R}$, we write $|a|$ for the absolute value of a, and, for a site $x = (x_1, \ldots, x_d) \in \mathbb{Z}^d$, we write $|x|$ for $\max(|x_1|, \ldots, |x_d|)$. For $n > 0$ and $x \in \mathbb{Z}^d$, let $Q_n(x) = \{y \in \mathbb{Z}^d \colon |y - x| \le n\}$ and $\partial Q_n(x) = \{y \in \mathbb{Z}^d \colon |y - x| = n\}$. We write Q_n for $Q_n(0)$ and ∂Q_n for $\partial Q_n(0)$.

For a positive integer r, we consider the torus $\mathbb{T}_r^d = (\{0, \ldots, r-1\}^d, \mathbb{E}_r^d)$ with the origin $0 = (0, \ldots, 0)$ and the edge set $\mathbb{E}_r^d = \big\{\{x, y\} \in \mathbb{Z}^d \times \mathbb{Z}^d \colon |x - y| = 1 (\text{mod } r)\big\}$. We denote the number of vertices in the torus or *volume* by $V = r^d$. For $p \in [0, 1]$, we consider the probability space $(\Omega_{\mathbb{T},p}, \mathcal{F}_{\mathbb{T},p}, \mathbb{P}_{\mathbb{T},p})$, where $\Omega_{\mathbb{T},p} = \{0, 1\}^{\mathbb{E}_r^d}$, $\mathcal{F}_{\mathbb{T},p}$ is the σ-field generated by the finite-dimensional cylinders of $\Omega_{\mathbb{T},p}$, and $\mathbb{P}_{\mathbb{T},p}$ is a product measure on $(\Omega_{\mathbb{T},p}, \mathcal{F}_{\mathbb{T},p})$, $\mathbb{P}_{\mathbb{T},p} = \prod_{e \in \mathbb{E}_r^d} \mu_e$, where μ_e is given by $\mu_e(\omega_e = 1) = 1 - \mu_e(\omega_e = 0) = p$, for vectors $(\omega_e)_{e \in \mathbb{E}_r^d} \in \Omega_{\mathbb{T},p}$. We write $\mathbb{E}_{\mathbb{T},p}$ for the expectation with respect to $\mathbb{P}_{\mathbb{T},p}$.

We further consider the hypercubic lattice $(\mathbb{Z}^d, \mathbb{E}^d)$, where the edge set is given by $\mathbb{E}^d = \big\{\{x,y\} \in \mathbb{Z}^d \times \mathbb{Z}^d \colon |x-y| = 1\big\}$. We often abuse notation and write $\mathbb{Z}^d$ for $(\mathbb{Z}^d, \mathbb{E}^d)$. For $p \in [0,1]$, we consider a probability space $(\Omega_{\mathbb{Z},p}, \mathcal{F}_{\mathbb{Z},p}, \mathbb{P}_{\mathbb{Z},p})$, where $\Omega_{\mathbb{Z},p} = \{0,1\}^{\mathbb{E}^d}$, $\mathcal{F}_{\mathbb{Z},p}$ is the σ-field generated by the finite-dimensional cylinders of $\Omega_{\mathbb{Z},p}$, and $\mathbb{P}_{\mathbb{Z},p}$ is the product measure on $(\Omega_{\mathbb{Z},p}, \mathcal{F}_{\mathbb{Z},p})$, $\mathbb{P}_{\mathbb{Z},p} = \prod_{e \in \mathbb{E}^d} \mu_e$, where μ_e is given by $\mu_e(\omega_e = 1) = 1 - \mu_e(\omega_e = 0) = p$, for vectors $(\omega_e)_{e \in \mathbb{E}^d} \in \Omega_{\mathbb{Z},p}$. In both settings, we say that an edge e is *open* or *occupied* if $\omega_e = 1$, and e is *closed* or *vacant* if $\omega_e = 0$.

The event that two sets of sites $\mathcal{K}_1, \mathcal{K}_2 \subset \mathbb{T}_r^d$ are connected by an open path is denoted by $\{\mathcal{K}_1 \leftrightarrow \mathcal{K}_2 \text{ in } \mathbb{T}_r^d\}$. We write $\mathcal{C}_{\mathbb{T}}(x)$ for the set of $y \in \mathbb{T}_r^d$ such that $x \leftrightarrow y$ in $\mathbb{T}_r^d$. Similarly, the event that two sets of sites $\mathcal{K}_1, \mathcal{K}_2 \subset \mathbb{Z}^d$ are connected by an open path is denoted by $\{\mathcal{K}_1 \leftrightarrow \mathcal{K}_2 \text{ in } \mathbb{Z}^d\}$, and the event that two sets of sites $\mathcal{K}_1, \mathcal{K}_2 \subset \mathbb{Z}^d$ are connected by an open path in Q_n is denoted by $\{\mathcal{K}_1 \leftrightarrow \mathcal{K}_2 \text{ in } Q_n\}$. The event that two sets of sites $\mathcal{K}_1, \mathcal{K}_2 \subset \mathbb{Z}^d$ are connected by an open path of length at most k is denoted by $\{\mathcal{K}_1 \xleftrightarrow{\leq k} \mathcal{K}_2 \text{ in } \mathbb{Z}^d\}$. We write $\mathcal{C}_{\mathbb{Z}}(x)$ for the set of $y \in \mathbb{Z}^d$ such that $x \leftrightarrow y$ in $\mathbb{Z}^d$. Finally, for $x, y \in \mathbb{T}_r^d$, we write $\tau_{\mathbb{T},p}(x,y) = \mathbb{P}_{\mathbb{T},p}(x \leftrightarrow y \text{ in } \mathbb{T}_r^d)$, and $\tau_{\mathbb{T},p}(x) = \tau_{\mathbb{T},p}(0,x)$, while, for $x, y \in \mathbb{Z}^d$, $\tau_{\mathbb{Z},p}(x,y) = \mathbb{P}_{\mathbb{Z},p}(x \leftrightarrow y \text{ in } \mathbb{Z}^d)$ and $\tau_{\mathbb{Z},p}(x) = \tau_{\mathbb{Z},p}(0,x)$.

For two nearest-neighbor vertices x and y in $\mathbb{T}_r^d$, we let $e_{x,y}$ be the unit vector $(y-x) \bmod(r)$. Let e_i be the i^{th} unit vector. We write $\langle e, f \rangle$ for the scalar product of vectors e and f. For a nearest-neighbor path $\pi = (x(1), \ldots, x(m))$ in $\mathbb{T}_r^d$, we define its *absolute winding* as

$$w(\pi) = \max_{i=1,\ldots,d} \left| \sum_{k=1}^{m-1} \langle e_{x(k),x(k+1)}, e_i \rangle \right|.$$

For a nearest-neighbor path $\pi = (x(1), \ldots, x(m))$ in $\mathbb{T}_r^d$, we define the subpath π_n by $\pi_n = (x(1), \ldots, x(n))$. We define the *maximal displacement* of π as

$$\delta(\pi) = \max_{n=1,\ldots,m} w(\pi_n). \tag{1.3}$$

This is an important notion for Proposition 2.1, which is in the heart of our proofs. We call $\pi = (x(1), \ldots, x(m))$ a *cycle* if $x(1) = x(m)$ and $x(i) \neq x(j)$ for all other $i \neq j$, i.e., π is a self-avoiding polygon. We say that a cycle π is *non-contractible* if $w(\pi) \neq 0$. (Note that $w(\pi) = 0 (\text{mod } r)$.) We further say that a cycle $\pi = (x(1), \ldots, x(m))$ is *long* if $\delta(\pi^{(n)}) \geq r/2$ for every $n = 1, \ldots, m$, where $\pi^{(n)} = (x(n), \ldots, x(m), x(1), \ldots, x(n))$ is the cycle π starting and ending at $x(n)$. Non-contractible cycles are examples of long cycles, but not every long cycle is non-contractible. Note further that if π is long then the reverse of π is also long.

For two functions g and h from a set $\mathcal{X}$ to $\mathbb{R}$, we write $g(z) \asymp h(z)$ to indicate that $g(z)/h(z)$ is bounded away from 0 and ∞, uniformly in $z \in \mathcal{X}$. Throughout this paper we write log for $\log_2$. All the constants (C_i) in the proofs are strictly positive and finite and depend only on the dimension, unless the dependence on other parameters is explicitly stated. Their exact values may be different from section to section.

We first give bounds on the probability that a vertex of the torus is in an open long cycle.

Theorem 1.1 (Expected number of vertices in long cycles)**.** *Assume (1.1). Let p_c be the critical threshold for bond percolation on $\mathbb{Z}^d$. For $x \in \mathbb{T}_r^d$,*

$$\mathbb{P}_{\mathbb{T},p_c}(x \textit{ is in an open long cycle}) \asymp V^{-2/3}. \tag{1.4}$$

Consequently,

$$\mathbb{E}_{\mathbb{T},p_c}\big[\#\{x \colon x \textit{ is in an open long cycle}\}\big] \asymp V^{1/3}. \tag{1.5}$$

In the next theorem we show that, with high probability, large open clusters of the torus may only contain long cycles of length of order $V^{1/3}$. Here, the *length* of a cycle is the number of edges it contains.

Theorem 1.2 (Cycles have length of order $V^{1/3}$). *Assume (1.1).*
(a) There exists $\delta = \delta(\varepsilon)$ *such that* $\delta \downarrow 0$ *as* $\varepsilon \downarrow 0$ *and*

$$\limsup_{\varepsilon\downarrow 0} \limsup_{V\to\infty} \mathbb{P}_{\mathbb{T},p_c}\Big(\exists x\colon |\mathcal{C}_{\mathbb{T}}(x)| > \delta V^{2/3},\ \mathcal{C}_{\mathbb{T}}(x) \text{ contains a long cycle of length } \leq \varepsilon V^{1/3}\Big) = 0. \tag{1.6}$$

(b) There exists a constant $C > 0$ *such that for all* $\varepsilon > 0$, *and* $r \geq 1$,

$$\mathbb{P}_{\mathbb{T},p_c}\Big(\exists \text{ a long cycle of length } \geq \varepsilon^{-1} V^{1/3}\Big) \leq C\varepsilon. \tag{1.7}$$

Theorem 1.2(b) follows rather directly from Theorem 1.1. We prove Theorem 1.2(a) and Theorem 1.1 at the same time. The proof of Theorem 1.2(a) consists of two main steps. We first rule out the possibility of the existence of long cycles with displacement shorter than $V^{1/6}$. In words, these are the cycles that do not wind around the torus too often. After that, we show that long cycles with few edges cannot have large displacement.

We next study the number of long cycles. We start by defining what this is. For a subgraph G of the torus, we define Y_G as the smallest k for which there exist edges $e_1, \ldots, e_k$ in G such that $G \setminus \{e_1, \ldots, e_k\}$ does not contain any long cycles. For $\delta > 0$, we define

$$Y_\delta = \sum Y_{\mathcal{C}} I(|\mathcal{C}| > \delta V^{2/3}),$$

where the sum is over all open clusters of the torus. We prove the following theorem:

Theorem 1.3 (A bound on the number of long cycles in large clusters). *Assume (1.1). There exists a finite constant* C *such that for all* r *and* $\delta > 0$,

$$\mathbb{E}_{\mathbb{T},p_c}[Y_\delta] \leq C/\delta.$$

In particular, the random variables Y_δ *are tight.*

Theorem 1.4 (Non-trivial existence of long cycles). *Assume (1.1).*
(a) There exists $c > 0$ *such that for all* $r \geq 1$,

$$\mathbb{P}_{\mathbb{T},p_c}\Big(\exists \text{ a long cycle of length } > cV^{1/3}\Big) > c.$$

(b) For any $\delta > 0$ *there exists* $c > 0$ *such that for all* $r \geq 1$,

$$\mathbb{P}_{\mathbb{T},p_c}(Y_\delta = 0) > c.$$

In other words, with positive probability uniformly in r, *there are no long cycles in clusters of size* $> \delta V^{2/3}$.

1.3 Related results on critical percolation

In this section, we state a few results that are interesting in their own right, for the ease of future reference. We start with a result that shows that short arms are unlikely:

Theorem 1.5 (Short arms are unlikely). *Assume (1.1). There exist constants* C, a *such that, for all* $\varepsilon > 0$,

$$\limsup_{n\to\infty} \mathbb{P}_{\mathbb{Z},p_c}\Big(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} \partial Q_n \mid 0 \leftrightarrow \partial Q_n\Big) \leq C \mathrm{e}^{-a/\sqrt{\varepsilon}}. \tag{1.8}$$

Theorem 1.6 (Connections inside balls)**.** *Assume (1.1). There exists $C < \infty$ such that*
(a) for all $n \geq 1$,

$$\sum_{x \in \partial Q_n} \mathbb{P}_{\mathbb{Z},p_c}(0 \leftrightarrow x \text{ in } Q_n) \leq C,$$

(b) for all $\varepsilon > 0$,

$$\limsup_{n\to\infty} \sum_{x \in \partial Q_n} \mathbb{P}_{\mathbb{Z},p_c}(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} x \text{ in } Q_n) \leq C\sqrt{\varepsilon}. \tag{1.9}$$

Theorem 1.7 (Torus two-point function)**.** *There exists a finite constant C such that for all $x \in \mathbb{T}_r^d$,*

$$\tau_{\mathbb{T},p_c}(x) \leq \tau_{\mathbb{Z},p_c}(x) + CV^{-2/3}, \tag{1.10}$$

and

$$\sup_{x \in \mathbb{T}_r^d} \mathbb{P}_{\mathbb{T},p_c}(0 \leftrightarrow x \text{ by a path with displacement } \geq r/6) \leq CV^{-2/3}. \tag{1.11}$$

1.4 Discussion

In this section, we compare our results to those for the Erdős-Rényi random graph (ERRG), as proved, for example, by Aldous in [2], and formulate some open problems. We refer to [2] for the extensive literature on the cycle structure of the ERRG. The ERRG is obtained by removing each edge of the complete graph K_n independently with probability p. On the ERRG, within the critical window $p = (1 + \lambda n^{-1/3})/n$, the number of cycles of large clusters is a random variable whose distribution can be explicitly characterized in terms of a Poisson random variable with a *random* parameter. This random parameter can be described as the area of the cluster exploration process. Since this parameter is a.s. bounded, in particular, each large cluster has an a.s. bounded number of cycles, and the probability that the i^{th} largest cluster does not contain any cycle is strictly positive. Since there is just a finite number of clusters of size at least $\delta n^{2/3}$ this immediately implies that the probability that there are no connected components of size at least $\delta n^{2/3}$ containing cycles is strictly positive. Further, all cycles have macroscopic length. Indeed, the largest connected components in the ERRG have diameter of the order $n^{1/3}$, and the length of cycles (when they exist) is also of the same order of magnitude. Cycles play a crucial role in describing the scaling limit of the largest critical clusters on the ERRG, as identified in [1]. Indeed, clusters locally look like trees, with cycles creating shortcuts between the different branches of the tree. Since cycles have a macroscopic length, these shortcuts are also macroscopic and thus the scaling limit of large critical clusters on the ERRG within the critical window is *not* a tree.

In this paper, we have proved similar properties for critical percolation on the high-dimensional torus, in order to provide yet another argument why the scaling limit of critical percolation on high-dimensional tori should be related to that for the critical ERRG.

We complete this section by formulating a few interesting extensions to our work. The first extension deals with the values of p within the so-called *scaling window*. The results in [14, 15], in conjunction with those in [5, 6], show that when $p = p_c(1 + \varepsilon)$, where $V^{1/3}|\varepsilon|$ remains uniformly bounded, the largest clusters obey similar scaling as for $p = p_c$. It would be of interest to extend our work to the full scaling window. The main difficulty therein lies in the extension of Theorems 1.5–1.6 to the relevant values of n and p. The second extension is to more general high-dimensional tori, for example, the n-cube as studied in [7]. Due to the simpler nature of the graph, it might be possible to prove stronger versions of our results. For example, is it possible to prove that the probability that there are no long cycles is strictly bounded away from 0 and 1?

Organization of this paper. This paper is organized as follows. In Section 2, we collect some preliminary results that we use in the proof. In Section 3, we prove Theorem 1.1. In Section 4, we prove Theorem 1.2. In Section 5, we prove Theorems 1.3 and 1.4. In Section 6, we prove Theorems 1.5 and 1.6, and in Section 7 we prove Theorem 1.7.

2 Preliminary results

In this section we collect some results that we use in the proofs.

2.1 Coupling clusters on the torus and the lattice

We say that two vertices x and y in $\mathbb{Z}^d$ are *r-equivalent* and write $x \overset{r}{\sim} y$, if $y = x + rz$ for some $z \in \mathbb{Z}^d$. We say that two edges $e_1 = \{x_1, y_1\}$ and $e_2 = \{x_2, y_2\}$ in $\mathbb{E}^d$ are *r-equivalent* ($e_1 \overset{r}{\sim} e_2$) if $x_1 \overset{r}{\sim} x_2$ and $y_1 \overset{r}{\sim} y_2$, or if $x_1 \overset{r}{\sim} y_2$ and $y_1 \overset{r}{\sim} x_2$.

Let $\Delta_n(x)$ be the event that the origin is connected to x in $\mathbb{T}_r^d$ by an open path π with $\delta(\pi) \le n$. We denote by $\mathcal{C}_{\mathbb{T}}(0; n)$ the set of all $x \in \mathbb{T}_r^d$ such that $\Delta_n(x)$ occurs. We also denote by $\mathcal{C}_{\mathbb{Z}}(0; n)$ the set of all $x \in \mathbb{Z}^d$ such that $0 \leftrightarrow x$ in Q_n. Let LC_n be the event that the origin is in an open long cycle $\pi = (x(1) = 0, \ldots, x(m))$ with $\delta(\pi) \le n$.

In the proof of Theorem 1.1, we need an extension of [14, Proposition 2.1]:

Proposition 2.1 (Coupling of clusters on torus and lattice)**.** *Consider bond percolation on $\mathbb{Z}^d$ and on $\mathbb{T}_r^d$ with parameter $p \in [0, 1]$. There exists a probability law $\mathbb{P}_{\mathbb{Z},\mathbb{T},p}$ on the joint space of percolation on $\mathbb{Z}^d$ and $\mathbb{T}_r^d$ such that, for all n and for all events E,*

$$\mathbb{P}_{\mathbb{Z},\mathbb{T},p}(\mathcal{C}_{\mathbb{T}}(0; n) \in E) = \mathbb{P}_{\mathbb{T},p}(\mathcal{C}_{\mathbb{T}}(0; n) \in E), \quad \mathbb{P}_{\mathbb{Z},\mathbb{T},p}(\mathcal{C}_{\mathbb{Z}}(0; n) \in E) = \mathbb{P}_{\mathbb{Z},p}(\mathcal{C}_{\mathbb{Z}}(0; n) \in E),$$

and $\mathbb{P}_{\mathbb{Z},\mathbb{T},p}$-almost surely, for all $x \in \mathbb{T}_r^d$,

$$\Delta_n(x) \subset \bigcup_{y \in Q_n, y \overset{r}{\sim} x} \{0 \leftrightarrow y \text{ in } Q_n\}.$$

Moreover, $\mathbb{P}_{\mathbb{Z},\mathbb{T},p}$-almost surely, (a) if $\mathcal{C}_{\mathbb{T}}(0; n) \neq \mathcal{C}_{\mathbb{T}}(0; n-1)$ then $0 \leftrightarrow \partial Q_n$ in $\mathbb{Z}^d$, and (b) if there exist $x, y \in \mathcal{C}_{\mathbb{T}}(0; n) \setminus \mathcal{C}_{\mathbb{T}}(0; n-1)$ (not necessarily distinct) such that the origin is connected to x and y in $\mathbb{T}_r^d$ by two disjoint open paths, then the origin is connected to ∂Q_n in $\mathbb{Z}^d$ by two disjoint open paths.

Finally, for $x \overset{r}{\sim} y$, and $\mathbb{P}_{\mathbb{Z},\mathbb{T},p}$-almost surely, the event

$$\{0 \leftrightarrow y \text{ in } Q_n\} \setminus \Delta_n(x)$$

implies that there exist distinct r-equivalent vertices v_1 and v_2 in Q_n and a vertex $z \in Q_n$ such that the following disjoint connections take place in Q_n:

$$\{0 \leftrightarrow z\} \circ \{z \leftrightarrow v_1\} \circ \{z \leftrightarrow v_2\} \circ \{v_1 \leftrightarrow y\}.$$

Proof. We prove the proposition by defining an exploration of $\mathcal{C}_{\mathbb{T}}(0; n)$ and $\mathcal{C}_{\mathbb{Z}}(0; n)$ simultaneously for all n from a percolation configuration on $\mathbb{Z}^d$. At each step k of the exploration, we will keep track of the following sets of edges: the active edges $A(k)$, the occupied edges $O(k)$, the vacant edges $V(k)$, the ghost edges $G(k)$, and the explored edges $E(k) = O(k) \cup V(k) \cup G(k)$.

We number the edges of $\mathbb{Z}^d$ in such a way that, for any two edges $e_1 = \{x, x'\}$ and $e_2 = \{y, y'\}$ in $\mathbb{Z}^d$, e_1 has a smaller number than e_2 if (a) $\max(|x|, |x'|) < \max(|y|, |y'|)$ or (b) $\max(|x|, |x'|) = \max(|y|, |y'|)$ and $\min(|x|, |x'|) < \min(|y|, |y'|)$. Thus, edges with small numbers are close to the origin.

We initiate the exploration by taking $A(0)$ to be the set of all edges that are neighbors of the origin, and $E(0) = \varnothing$. Assume that $A(k)$, $O(k)$, $V(k)$ and $G(k)$ are defined. If $A(k) = \varnothing$, then we stop the exploration. Otherwise, we take the edge $e \in A(k)$ with the smallest number. We define $G(k+1) = G(k) \cup \{f \colon f \neq e, f \overset{r}{\sim} e\}$. To make the other updates, we consider two cases: (a) if e is open, we define $O(k+1) = O(k) \cup \{e\}$, $V(k+1) = V(k)$, and $A(k+1) = A(k) \cup \{f \colon f \sim e\} \setminus E(k+1)$, where $e \sim f$ means that e and f share an end-vertex; (b) if e is closed, we define $O(k+1) = O(k)$, $V(k+1) = V(k) \cup \{e\}$, and $A(k+1) = A(k) \setminus E(k+1)$.

The exploration eventually stops at some step $T+1$, which is the first step T such that $A(T+1) = \varnothing$. Note that T is at most the number of edges in $\mathbb{T}_r^d$, i.e., $T \leq dV$. We call the above procedure the $\mathbb{T}$-*exploration.* As a result of the $\mathbb{T}$-exploration, we obtain an embedding $O(T)$ of the edge set of $\mathcal{C}_{\mathbb{T}}(0)$ into $\mathbb{Z}^d$. However, because of the particular choice of the numbering of the edges of $\mathbb{Z}^d$ used for the exploration, which is such that if $e_1 < e_2$, then e_1 is closer to the origin that e_2, we can say much more about the structure of the embedding. Indeed, let $T_n + 1$ be the first time an edge from Q_n^c is explored. Then $O(T_n)$ is an embedding of the edge set of $\mathcal{C}_{\mathbb{T}}(0; n)$ into Q_n.

The exploration of $\mathcal{C}_{\mathbb{Z}}(0)$ is similar to the $\mathbb{T}$-exploration, with the only difference that $G(k)$ is taken to be the empty set for all k. This exploration may terminate at some finite time, in which case $\mathcal{C}_{\mathbb{Z}}(0)$ is finite, or continue forever, which corresponds to the case of infinite $\mathcal{C}_{\mathbb{Z}}(0)$.

Most of the properties of the coupling (defined in the statement of the proposition) are immediate from the construction.

To see that the last property holds, we assume that the event $0 \leftrightarrow y$ in Q_n holds for some $x \overset{r}{\sim} y$, while $\Delta_n(x)$ does not hold. We fix an open path π from 0 to y in Q_n. Since $\Delta_n(x)$ does not occur, there exists an edge e on this path, such that $e \in G(T_n)$. Let e be the last edge on π from $G(T_n)$. By construction, there exists an edge $f \in V(T_n) \subset Q_n$ such that $f \neq e$, $f \overset{r}{\sim} e$, and the origin is connected to one of the end vertices of f by an open path inside $O(T_n)$. We denote this vertex by v_2, and let v_1 be the end-vertex of e which is r-equivalent to v_2. Note that none of the edges in the part of π from e to y is in $O(T_n)$. Indeed, let $\{g_1, \ldots, g_s\} \subset Q_n$ be the edges in the part of π from e to y, and assume that $g_i \in O(T_n)$ for some $i \in \{1, \ldots, s\}$. Since all the g_j's are open on $\mathbb{Z}^d$, are in Q_n, and, by the definition of e, none of them is a ghost edge, we conclude that the edges g_j, $j \geq i$ are also in $O(T_n)$. This is in contradiction to the assumption that $\Delta_n(x)$ does not occur. Therefore, we conclude that the event $\{0 \leftrightarrow v_1, v_2 \text{ in } Q_n\} \circ \{v_1 \leftrightarrow y \text{ in } Q_n\}$ occurs. We finish the proof by observing that if 0 is connected to v_1 and v_2 by open paths in Q_n, then there exists $z \in Q_n$ such that the following disjoint open paths exist in Q_n: $\{0 \leftrightarrow z\} \circ \{z \leftrightarrow v_1\} \circ \{z \leftrightarrow v_2\}$. □

The following corollary is used in the proof of Theorem 1.1:

Corollary 2.1. *There exists a probability law on the joint space of percolation on $\mathbb{Z}^d$ and $\mathbb{T}_r^d$ such that if the origin is in an open long cycle $\pi = (x(1) = 0, \ldots, x(m))$ in $\mathbb{T}_r^d$ and the event* LC_{3n} *does not occur, then there exist two disjoint open paths from the origin to ∂Q_n in $\mathbb{Z}^d$.*

Proof. We use the coupling from Proposition 2.1. We define the reverse of a path $\bar{\pi} = (x(1), \ldots, x(m))$ as $\bar{\pi}^{-1} = (x(m), \ldots, x(1))$. Note that for any path $\bar{\pi}$,

$$\delta(\bar{\pi}^{-1}) \leq 2\delta(\bar{\pi}).$$

Indeed, consider the path $\bar{\pi}$ as the concatenation of two paths $\bar{\pi}_1 = (x(1), \ldots, x(k))$ and $\bar{\pi}_2 = (x(k), \ldots, x(m))$ such that $\delta(\bar{\pi}^{-1}) = \delta(\bar{\pi}_2^{-1}) = w(\bar{\pi}_2)$. It follows from the triangle inequality that $w(\bar{\pi}_2) \leq w(\bar{\pi}) + w(\bar{\pi}_1)$. Moreover, by the definition of the maximal displacement of $\bar{\pi}$, we have the bounds $w(\bar{\pi}) \leq \delta(\bar{\pi})$ and $w(\bar{\pi}_1) \leq \delta(\bar{\pi})$.

Let π be an open long cycle starting at the origin. If LC_{3n} does not occur, then $\delta(\pi) > 3n$. Moreover, there exists $x = x(k) \in \pi$ such that for any pair of disjoint open paths π_1 and π_2 from 0 to x and from x

to 0, respectively, which form a long cycle, their displacements satisfy $\delta(\pi_1) \geq n$ and $\delta(\pi_2) \geq 2n$. Indeed, one can take x on the boundary of $\mathcal{C}_{\mathbb{T}}(0;n)$. (Mind that $\delta(\pi_1) + \delta(\pi_2) \geq \delta(\pi)$.) Since $\delta(\pi_2) \geq 2n$, by the above property of the displacement of the reverse path, we have $\delta(\pi_2^{-1}) \geq \delta(\pi_2)/2 \geq n$. The result now follows from Proposition 2.1(b). $\square$

From now on, we only consider the probability measure $\mathbb{P}_{\mathbb{Z},\mathbb{T},p}$ defined in Proposition 2.1 and

in the remainder of the paper, we write $\mathbb{P}_p$ *for* $\mathbb{P}_{\mathbb{Z},\mathbb{T},p}$.

In particular, $\mathbb{P}_p(E) = \mathbb{P}_{\mathbb{Z},p}(E)$ for $E \in \mathcal{F}_{\mathbb{Z},p}$, and $\mathbb{P}_p(E) = \mathbb{P}_{\mathbb{T},p}(E)$ for $E \in \mathcal{F}_{\mathbb{T},p}$ (see Section 2.1 for notation), and we always assume without mentioning the coupling from Proposition 2.1 when we consider events from $\mathcal{F}_{\mathbb{Z},p}$ and $\mathcal{F}_{\mathbb{T},p}$ simultaneously.

2.2 Previous results

In the next theorem, we summarize a number of results on high-dimensional percolation on $\mathbb{Z}^d$ that we will often use in the proofs in this paper.

Theorem 2.1 (Critical behavior of high-dimensional percolation). *Assume (1.1). Then, there exist* $c > 0$ *and* $C < \infty$ *such that for all positive integers* n*,*

(i)

$$cn^{-2} \leq \mathbb{P}_{p_c}\left(0 \leftrightarrow \partial Q_n \text{ in } \mathbb{Z}^d\right) \leq Cn^{-2}, \tag{2.1}$$

(ii)

$$cn^2 \leq \sum_{x \in Q_n} \tau_{\mathbb{Z},p_c}(x) \leq Cn^2, \tag{2.2}$$

and for any given $z \in \mathbb{Z}^d$ *and a positive integer* r *with* $r \leq n$*,*

$$\frac{cn^2}{r^d} \leq \sum_{x \in Q_n, x \overset{r}{\sim} z, |x| \geq r/4} \tau_{\mathbb{Z},p_c}(x) \leq \frac{Cn^2}{r^d}. \tag{2.3}$$

(iii) For any $z \in \mathbb{Z}^d$*,*

$$\sum_{x,y \in \mathbb{Z}^d} \tau_{\mathbb{Z},p_c}(0,x)\tau_{\mathbb{Z},p_c}(x,y)\tau_{\mathbb{Z},p_c}(y,z) \leq C|z|^{6-d}. \tag{2.4}$$

Proof. The first statement is [19, Theorem 1]. Statements (ii) and (iii) easily follow from (1.1). $\square$

The next theorem gives an upper bound on $\mathbb{E}_{\mathbb{T},p_c}|\mathcal{C}(0)|$, which is used often in the proofs:

Theorem 2.2 (Expected critical cluster size on torus). *There exists a finite constant* C *such that for all* $r \geq 1$*,*

$$\mathbb{E}_{\mathbb{T},p_c}|\mathcal{C}(0)| \leq CV^{1/3}. \tag{2.5}$$

Proof. The statement follows from [14, (1.6)] and [6, Theorem 1.6(iii)]. Alternatively, it follows from (1.10) and (2.2), where we note that the proof of (1.10) in Section 7 only relies on (2.1) and Proposition 2.1. $\square$

3 Proof of Theorem 1.1

In this section, we prove Theorem 1.1. We give the proof of the upper bound in Proposition 3.1 and the proof of the lower bound in Proposition 3.2. The results of these propositions are more general than the result of Theorem 1.1, and also give bounds on the probability that the origin is in a long cycle with small displacement. In particular, the upper bound on the probability of such an event will be used in the proof of Theorem 1.2 to show that large open clusters do not contain long cycles with few edges.

Recall the definitions of $\Delta_n(x)$ and LC_n from Section 2.1. We prove the following bounds:

Proposition 3.1 (Upper bound on long cycles)**.** *Assume (1.1). There exists a constant C such that*

$$\mathbb{P}_{p_c}(0 \textit{ is in an open long cycle}) \le CV^{-2/3},$$

and

$$\mathbb{P}_{p_c}(\mathrm{LC}_n) \le Cn^2/V.$$

Proof. Let A be the event that the origin in an open long cycle and the event $\mathrm{LC}_{3V^{1/6}}$ does not occur. We split

$$\mathbb{P}_{p_c}(0 \text{ is in an open long cycle}) = \mathbb{P}_{p_c}(A) + \mathbb{P}_{p_c}(\mathrm{LC}_{3V^{1/6}}). \tag{3.1}$$

We first estimate the probability of A. By Corollary 2.1, the event A implies the existence of two disjoint paths from 0 to $\partial Q_{V^{1/6}}$ in $\mathbb{Z}^d$. This observation and the BK inequality (see, e.g., [10, Theorem 2.12]) imply

$$\mathbb{P}_{p_c}(A) \le \mathbb{P}_{p_c}\left(0 \longleftrightarrow \partial Q_{V^{1/6}} \text{ in } \mathbb{Z}^d\right)^2 \le C_1 V^{-2/3},$$

where the last inequality follows from (2.1).

We now bound the probability of the event $\mathrm{LC}_{3V^{1/6}}$. The event $\mathrm{LC}_{3V^{1/6}}$ implies that there exists a vertex $x \in \partial Q_{\lfloor r/2 \rfloor}$ such that (a) there are two disjoint paths π_1 and π_2 from 0 to x in $\mathbb{T}_r^d$, (b) the path π_1 is contained inside $Q_{\lfloor r/2 \rfloor}$, and (c) $\delta(\pi_2) \le 3V^{1/6}$. We use the BK inequality to bound the probability of the event $\mathrm{LC}_{3V^{1/6}}$ by

$$\sum_{x \in \partial Q_{\lfloor r/2 \rfloor}} \mathbb{P}_{p_c}\left(0 \leftrightarrow x \text{ in } Q_{\lfloor r/2 \rfloor}\right) \mathbb{P}_{p_c}\left(\Delta_{3V^{1/6}}(x)\right).$$

It follows from Proposition 2.1 that if $\Delta_{3V^{1/6}}(x)$ occurs, then there exists $y \in \mathbb{Z}^d$ with $y \overset{r}{\sim} x$ and $|y| \le V^{1/6}$ such that $0 \leftrightarrow y$ in $Q_{3V^{1/6}}$. By (2.3) and the fact that $|y| \ge r/2$,

$$\mathbb{P}_{p_c}\left(\Delta_{3V^{1/6}}(x)\right) \le \sum_{y \in Q_{3V^{1/6}}, y \overset{r}{\sim} x} \mathbb{P}_{p_c}(0 \leftrightarrow y \text{ in } \mathbb{Z}^d) \le C_2 V^{-2/3}, \tag{3.2}$$

where the constant C_2 does not depend on x or r. Finally, Theorem 1.6(a) implies

$$\sum_{x \in \partial Q_{\lfloor r/2 \rfloor}} \mathbb{P}_{p_c}\left(0 \leftrightarrow x \text{ in } Q_{\lfloor r/2 \rfloor}\right) \le C_3. \tag{3.3}$$

Therefore, we arrive at

$$\mathbb{P}_{p_c}(\mathrm{LC}_{3V^{1/6}}) \le C_4 V^{-2/3}.$$

This completes the proof of the first statement of Proposition 3.1. The proof of the second statement is similar to the proof of the bound for $\mathbb{P}_{p_c}(\mathrm{LC}_{3V^{1/6}})$, now for n instead of $3V^{1/6}$. In particular, (3.2) is replaced by

$$\mathbb{P}_{p_c}\left(\Delta_n(x)\right) \le \sum_{y \in Q_n, y \overset{r}{\sim} x} \mathbb{P}_{p_c}(0 \leftrightarrow y \text{ in } \mathbb{Z}^d) \le C_5 n^2/V.$$

We omit further details. □

Proposition 3.2 (Lower-bound on long cycles)**.** *Assume (1.1). There exist positive constants* ε_0 *and* c *such that*

$$\mathbb{P}_{p_c}(0 \text{ is in an open long cycle}) \geq cV^{-2/3},$$

and, for any $n \in [\varepsilon_0^{-1} r, \varepsilon_0 V^{1/6}]$,

$$\mathbb{P}_{p_c}(\mathrm{LC}_n) \geq cn^2/V.$$

Proof. The first statement immediately follows from the second one for $n = \varepsilon_0 V^{1/6}$. We now prove the second statement. Let K be a large positive integer. Let ε and α be small positive numbers. The precise choice of these numbers will be made later in the proof. Let n be an integer in $[\alpha^{-1}r, \varepsilon V^{1/6}]$.

First of all, note that it suffices to prove the result for $r > 4K$. Indeed, once we fix K, the result for $r \in [1, 4K]$ will follow by adjusting the constant c. Therefore, throughout the proof we assume that $r > 4K$. The proof consists of several steps.

Step 1. For any $x \in \mathbb{Z}^d$, let $u = u(x)$ be the vertex in $\mathbb{Z}^d$ with coordinates $(x_1, \ldots, x_{d-1}, x_d + K)$. In particular, $u(x) \in \partial Q_K(x)$. For $x \in \mathbb{Z}^d$, consider the event

$$A_x = A_x(n, K) = \{0 \leftrightarrow u(x) \text{ in } Q_n\}.$$

Let

$$N(A) = \left| \{x \in \mathbb{Z}^d \colon x \overset{r}{\sim} 0,\ x \neq 0,\ A_x \text{ occurs}\} \right|.$$

We use the second moment method to show that

$$\mathbb{P}_{p_c}(N(A) \neq 0) \geq cn^2/V,$$

for some constant c which does not depend on r, n, ε or K.

We first show that

$$\mathbb{E}_{p_c} N(A) \geq C_6 n^2/V. \tag{3.4}$$

We take $\alpha \in (0, 1/2)$ and write

$$\begin{aligned} \mathbb{E}_{p_c} N(A) &\geq \sum_{x \in Q_{\alpha n} \setminus \{0\}, x \overset{r}{\sim} 0} \mathbb{P}_{p_c}(0 \leftrightarrow u(x) \text{ in } Q_n) \\ &= \sum_{x \in Q_{\alpha n} \setminus \{0\}, x \overset{r}{\sim} 0} \mathbb{P}_{p_c}(0 \leftrightarrow u(x)) - \sum_{x \in Q_{\alpha n} \setminus \{0\}, x \overset{r}{\sim} 0} \mathbb{P}_{p_c}(\{0 \leftrightarrow u(x)\} \setminus \{0 \leftrightarrow u(x) \text{ in } Q_n\}). \end{aligned}$$

Since $r \leq \alpha n$ and $r > 4K$, the first sum is bounded from below by $C_7(\alpha n)^2/V$ by (1.1) and (2.2). We use the BK inequality, Theorem 1.6(a), (2.3), (1.1) and the assumption $r > 4K$, to bound the second sum from above by

$$\begin{aligned} &\sum_{x \in Q_{\alpha n} \setminus \{0\}, x \overset{r}{\sim} 0} \sum_{y \in \partial Q_n} \mathbb{P}_{p_c}\Big(\{0 \leftrightarrow y \text{ in } Q_n\} \circ \{y \leftrightarrow u(x)\}\Big) \\ &\leq \sum_{x \in Q_{\alpha n} \setminus \{0\}, x \overset{r}{\sim} 0} \sum_{y \in \partial Q_n} \mathbb{P}_{p_c}(0 \leftrightarrow y \text{ in } Q_n) \mathbb{P}_{p_c}(y \leftrightarrow u(x)) \leq C_8(\alpha^d n^2)/V. \end{aligned} \tag{3.5}$$

The desired lower bound on $\mathbb{E}_{p_c} N(A)$ follows by taking α small enough. From this moment on, α remains unchanged.

Next, we bound the second moment of $N(A)$. Let $\sum'$ be the sum over all distinct $x, y \in Q_n$ such that $x, y \neq 0$ and $x, y \overset{r}{\sim} 0$. We obtain

$$\mathbb{E}_{p_c} N(A)^2 \leq \mathbb{E}_{p_c} N(A) + \sum\nolimits' \mathbb{P}_{p_c}\left(0 \leftrightarrow u(x) \text{ in } Q_n, 0 \leftrightarrow u(y) \text{ in } Q_n\right).$$

Note that if $0 \leftrightarrow u(x)$ in Q_n and $0 \leftrightarrow u(y)$ in Q_n, then there exists $z \in Q_n$ such that the following open paths occur disjointly: $0 \leftrightarrow z$, $z \leftrightarrow u(x)$ and $z \leftrightarrow u(y)$. Therefore, the BK inequality implies

$$\mathbb{E}_{p_c} N(A)^2 \leq \mathbb{E}_{p_c} N(A) + {\sum}' \sum_{z \in Q_n} \tau_{\mathbb{Z},p_c}(z) \tau_{\mathbb{Z},p_c}(z, u(x)) \tau_{\mathbb{Z},p_c}(z, u(y)).$$

By translation invariance and the fact that $u(x) - z = u(x - z)$, the product $\tau_{\mathbb{Z},p_c}(z)\tau_{\mathbb{Z},p_c}(z, u(x))\tau_{\mathbb{Z},p_c}(z, u(y))$ equals $\tau_{\mathbb{Z},p_c}(-z)\tau_{\mathbb{Z},p_c}(u(x - z))\tau_{\mathbb{Z},p_c}(u(y - z))$. Let ${\sum}''$ be the sum over all pairwise distinct $x, y, z \in Q_{2n}$ such that $x \overset{r}{\sim} y \overset{r}{\sim} z$. We obtain

$$\mathbb{E}_{p_c} N(A)^2 \leq \mathbb{E}_{p_c} N(A) + {\sum}'' \tau_{\mathbb{Z},p_c}(u(x)) \tau_{\mathbb{Z},p_c}(u(y)) \tau_{\mathbb{Z},p_c}(z).$$

Since x, y and z are distinct and r-equivalent, at least two of them are at distance at least $r/2$ from the origin. Therefore, the above sum is at most

$$\mathbb{E}_{p_c} N(A) + \sum_{x \overset{r}{\sim} y \overset{r}{\sim} z; x,y,z \in Q_{2n}; |x|,|y| \geq r/2} \left[2\tau_{\mathbb{Z},p_c}(u(x)) \tau_{\mathbb{Z},p_c}(y) \tau_{\mathbb{Z},p_c}(u(z)) + \tau_{\mathbb{Z},p_c}(u(x)) \tau_{\mathbb{Z},p_c}(u(y)) \tau_{\mathbb{Z},p_c}(z) \right].$$

Remember that we assume that $r > 4K$. Therefore, $|u(x)| \geq r/4$ when $|x| \geq r/2$. We can use (2.2), (2.3) and the fact that $r \leq n \leq \varepsilon V^{1/6}$ to bound the above sum by

$$\mathbb{E}_{p_c} N(A) + C_9 (n^2)(n^2/V)^2 \leq \mathbb{E}_{p_c} N(A) + C_9 (\varepsilon^4 V^{-1/3}) n^2/V \leq 2\mathbb{E}_{p_c} N(A), \tag{3.6}$$

when ε is sufficiently small. A second moment estimate, using (3.4) and (3.6), yields

$$\mathbb{P}_{p_c}(N(A) \neq 0) \geq \frac{(\mathbb{E}_{p_c} N(A))^2}{\mathbb{E}_{p_c} N(A)^2} \geq \frac{C_6 n^2}{2V}. \tag{3.7}$$

Step 2. Consider the event

$$E = E(r, n, K) = \{0 \leftrightarrow u(0) \text{ in } \mathbb{T}_r^d \text{ by an open path } \pi \text{ with } w(\pi) \geq r/2 \text{ and } \delta(\pi) \leq n\}.$$

In other words, the event E implies that 0 is connected to $u(0)$ in $\mathbb{T}_r^d$ by a path which, considered as a continuous path in the continuous torus, cannot be contracted into a path from 0 to $u(0)$ in Q_K. Moreover, this path does not wind around the torus too often. We show that for small enough ε and large enough K,

$$\mathbb{P}_{p_c}(E) \geq \frac{1}{2} \mathbb{P}_{p_c}(N(A) \neq 0).$$

Since $\mathbb{P}_{p_c}(E) \geq \mathbb{P}_{p_c}(N(A) \neq 0) - \mathbb{P}_{p_c}(\{N(A) \neq 0\} \setminus E)$, we should show that

$$\mathbb{P}_{p_c}(\{N(A) \neq 0\} \setminus E) \leq \frac{1}{2} \mathbb{P}_{p_c}(N(A) \neq 0), \tag{3.8}$$

when ε is chosen small enough, and K large enough.

If $N(A) \neq 0$ and E does not occur, then according to Proposition 2.1, there exist $x \in Q_n$ with $x \overset{r}{\sim} 0$ and $x \neq 0$, a vertex $z \in Q_n$, and distinct vertices v_1 and v_2 in Q_n with $v_1 \overset{r}{\sim} v_2$ such that the following disjoint connections take place in Q_n:

$$\{0 \leftrightarrow z\} \circ \{z \leftrightarrow v_1\} \circ \{z \leftrightarrow v_2\} \circ \{v_1 \leftrightarrow u(x)\}.$$

Therefore, the probability of the event $\{N(A) \neq 0\} \setminus E$ is bounded from above by

$$\sum_{x \in Q_n, x \overset{r}{\sim} 0, x \neq 0} \; \sum_{v_1, v_2 \in Q_n, v_1 \overset{r}{\sim} v_2, v_1 \neq v_2} \; \sum_{z \in Q_n} \tau_{\mathbb{Z},p_c}(z) \tau_{\mathbb{Z},p_c}(z, v_1) \tau_{\mathbb{Z},p_c}(z, v_2) \tau_{\mathbb{Z},p_c}(v_1, u(x)). \tag{3.9}$$

Note that, since v_1 and v_2 are distinct and r-equivalent, either $|v_1 - z| \geq r/2$ or $|v_2 - z| \geq r/2$. Assume first that $|v_2 - z| \geq r/2$. It follows from (2.3) that for any v_1 and z,

$$\sum_{v_2 \in Q_n : v_2 \overset{r}{\sim} v_1, v_2 \neq v_1, |v_2 - z| \geq r/2} \tau_{\mathbb{Z},p_c}(z, v_2) \leq C_{10} n^2 / V,$$

where C_{10} does not depend on n, v_1 or z. Further, it follows from (2.4) that

$$\sum_{z \in Q_n} \sum_{v_1 \in Q_n} \sum_{x \in Q_n, x \overset{r}{\sim} 0, x \neq 0} \tau_{\mathbb{Z},p_c}(z) \tau_{\mathbb{Z},p_c}(z, v_1) \tau_{\mathbb{Z},p_c}(v_1, u(x)) \leq \sum_{x \in Q_n, x \overset{r}{\sim} 0, x \neq 0} C_{11} |u(x)|^{6-d} \leq C_{12} \varepsilon^6, \tag{3.10}$$

where in the last inequality we used the assumption that $r > 4K$, implying that $|u(x)| \geq |x|/2$ for $x \overset{r}{\sim} 0$, and the fact that $n \leq \varepsilon V^{1/6}$. Therefore, for v_2 and z with $|v_2 - z| \geq r/2$, the sum (3.9) is, uniformly in K, bounded from above by

$$C_{10} C_{12} \varepsilon^6 n^2 / V.$$

Next, consider the sum (3.9) in the case $|v_1 - z| \geq r/2$. By translation invariance, the sum (3.9) equals

$$\sum_{x \in Q_n, x \overset{r}{\sim} 0, x \neq 0} \; \sum_{v_1, v_2 \in Q_n, v_1 \overset{r}{\sim} v_2, v_1 \neq v_2} \; \sum_{z \in Q_n} \tau_{\mathbb{Z},p_c}(z) \tau_{\mathbb{Z},p_c}(z, v_1) \tau_{\mathbb{Z},p_c}(z, v_2) \tau_{\mathbb{Z},p_c}(v_2, u(x) + (v_2 - v_1)).$$

By the definition of $u(x)$, the translation of $u(x)$ by $(v_2 - v_1)$ equals $u(x + (v_2 - v_1))$. Since $v_1 \overset{r}{\sim} v_2$, the translation of x (r-equivalent to 0) by $(v_2 - v_1)$ is still r-equivalent to 0. Note that $x + (v_2 - v_1) \in Q_{2n}$ and it is also possible that $x + (v_2 - v_1) = 0$. These observations imply that the above sum is bounded from above by

$$\sum_{x \in Q_{2n}, x \overset{r}{\sim} 0} \; \sum_{v_1, v_2 \in Q_n, v_1 \overset{r}{\sim} v_2, v_1 \neq v_2} \; \sum_{z \in Q_n} \tau_{\mathbb{Z},p_c}(z) \tau_{\mathbb{Z},p_c}(z, v_1) \tau_{\mathbb{Z},p_c}(z, v_2) \tau_{\mathbb{Z},p_c}(v_2, u(x)).$$

Since we only consider the above sum in the case z and v_1 satisfy $|v_1 - z| \geq r/2$, we obtain as before that

$$\sum_{v_1 \in Q_n, v_1 \overset{r}{\sim} v_2, v_1 \neq v_2, |v_1 - z| \geq r/2} \tau_{\mathbb{Z},p_c}(z, v_1) \leq C_{10} n^2 / V.$$

It remains to bound the sum

$$\sum_{z \in Q_n} \sum_{v_2 \in Q_n} \sum_{x \in Q_{2n}, x \overset{r}{\sim} 0} \tau_{\mathbb{Z},p_c}(z) \tau_{\mathbb{Z},p_c}(z, v_2) \tau_{\mathbb{Z},p_c}(v_2, u(x)).$$

There are two cases depending on whether $x \neq 0$ or $x = 0$. The case $x \neq 0$ can be considered similarly to (3.10), so the above sum is bounded from above by $C_{13} \varepsilon^6$ in this case (the constant changes, since we now sum over $x \in Q_{2n}$ instead of $x \in Q_n$).

It remains to consider the case $x = 0$. In this case $|u(x)| = K$, and we only need to bound the sum over $z, v_2 \in \mathbb{Z}^d$. This sum is bounded from above by $C_{14} K^{6-d}$ by (2.4), where C_{14} is independent of K. Therefore, for v_1 and z with $|v_1 - z| \geq r/2$, the sum (3.9) is bounded from above by

$$C_{10} (C_{13} \varepsilon^6 + C_{14} K^{6-d}) n^2 / V.$$

Now take K large and ε small so that (3.8) holds. We obtain from (3.7) and (3.8) that

$$\mathbb{P}_{p_c}(E) \geq C_6 n^2/(4V).$$

Step 3. We now show that there exists $C_{15} = C_{15}(K)$ such that

$$\mathbb{P}_{p_c}(\mathrm{LC}_{n+3K}) \geq C_{15}(K)\mathbb{P}_{p_c}(E).$$

This follows from a local modification argument as follows. Note that if E occurs, there exist z and z' on ∂Q_K such that z is connected to z' by a path in $\mathbb{T}_r^d \setminus Q_K$ which can not be contracted into a path from z to z' inside Q_K and, at the same time, does not wind around the torus too often. More precisely, this path π satisfies $w(\pi) \geq r/2$ and $\delta(\pi) \leq n + K$. We can therefore modify the configuration of bonds inside Q_K to make sure that $\{0 \leftrightarrow z \text{ in } Q_K\} \circ \{0 \leftrightarrow z' \text{ in } Q_K\}$, which implies that the origin is in a long cycle π' with $\delta(\pi') \leq n + 3K$. Since there are only finitely many edges in Q_K, the above inequality follows.

We can now complete the proof of Proposition 3.2. We pick α so that (3.4) holds for all $r > 4K$. We then pick ε and K to satisfy (3.6) and (3.8). It follows from Steps 2 and 3 of the proof that for all $r > 4K$ and $n \in [\alpha^{-1}r + 3K, \varepsilon V^{1/6}]$,

$$\mathbb{P}_{p_c}(\mathrm{LC}_n) \geq C_{15}\frac{C_6(n-3K)^2}{4V} \geq C_{16}n^2/V.$$

Finally, we adjust the constant C_{16} so that the result remained valid for $r \leq 4K$. □

4 Proof of Theorem 1.2

We start by proving Theorem 1.2(b), which is a simple consequence of Theorem 1.1. Indeed, when there exists a long cycle of length at least $\varepsilon^{-1}V^{1/3}$, then the number of vertices in long cycles is at least $\varepsilon^{-1}V^{1/3}$. Denote the number of vertices in long cycles by M. Then, by the Markov inequality and Theorem 1.1,

$$\begin{aligned} \mathbb{P}_{p_c}\Big(\exists \text{ a long cycle of length } \geq \varepsilon^{-1}V^{1/3}\Big) &\leq \mathbb{P}_{p_c}(M \geq \varepsilon^{-1}V^{1/3}) \\ &\leq \varepsilon V^{-1/3}\mathbb{E}_{p_c}[M] \leq C_1\varepsilon, \end{aligned} \tag{4.1}$$

by Theorem 1.1. In the remainder of this section, we prove Theorem 1.2(a).

In the first lemma of this section, we provide sufficient conditions for non-existence (with high probability) of long cycles with few edges contained in large open clusters. A more intuitive sufficient condition will be given later in Lemma 4.2.

Lemma 4.1. *Assume (1.1). Let ε be a positive number, and let k and n be positive numbers (possibly depending on ε and V) satisfying the following assumption:*

$$\limsup_{\varepsilon\downarrow 0}\limsup_{V\to\infty}\max\left\{\frac{n^2}{V^{1/3}}, V^{1/3}\mathbb{P}_{\mathbb{Z},p_c}\Big(0 \overset{\leq k}{\longleftrightarrow} \partial Q_n\Big)\right\} = 0.$$

Let $\delta = \delta(\varepsilon)$ be such that

$$\limsup_{\varepsilon\downarrow 0}\frac{1}{\delta(\varepsilon)}\limsup_{V\to\infty}\max\left\{\frac{n^2}{V^{1/3}}, V^{1/3}\mathbb{P}_{\mathbb{Z},p_c}\Big(0 \overset{\leq k}{\longleftrightarrow} \partial Q_n\Big)\right\} = 0.$$

Then,

$$\limsup_{\varepsilon\downarrow 0}\limsup_{V\to\infty}\mathbb{P}_{\mathbb{T},p_c}\Big(\exists x\colon |\mathcal{C}_{\mathbb{T}}(x)| > \delta V^{2/3} \textit{ and } \mathcal{C}_{\mathbb{T}}(x) \textit{ contains a long cycle of length } \leq k\Big) = 0.$$

Remark 1. For us, the most interesting choice of δ in Lemma 4.1 is such that $\delta = \delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$. For such δ, Lemma 4.1 gives sufficient conditions for Theorem 1.2. It is possible to make such a choice because of the assumption of Lemma 4.1.

Proof. We first give an upper bound on the probability that the origin is in a long cycle of length at most k. Let $\mathcal{E}_k$ denote this event. Recall that LC_{3n} is the event that the origin is in a long cycle π with $\delta(\pi) \leq 3n$. It follows from Theorem 1.1 that $\mathbb{P}_{p_c}(\mathrm{LC}_{3n}) \leq C_2 n^2/V$. It follows from Corollary 2.1 that the event $\mathcal{E}_k \setminus \mathrm{LC}_{3n}$ implies the existence of two disjoint paths of length at most k from the origin to ∂Q_n in $\mathbb{Z}^d$. These observations imply

$$\mathbb{P}_{p_c}(\mathcal{E}_k) \leq \mathbb{P}_{p_c}(\mathrm{LC}_{3n}) + \mathbb{P}_{p_c}(\mathcal{E}_k \setminus \mathrm{LC}_{3n}) \leq C_2 n^2/V + \mathbb{P}_{p_c}\left(0 \overset{\leq k}{\longleftrightarrow} \partial Q_n\right)^2, \tag{4.2}$$

where in the last bound we also used the BK inequality.

It is now easy to finish the proof of Lemma 4.1. Indeed, by the Markov inequality and translation invariance,

$$\begin{aligned}&\mathbb{P}_{p_c}\left(\exists x\colon |\mathcal{C}_{\mathbb{T}}(x)| > \delta V^{2/3} \text{ and } \mathcal{C}_{\mathbb{T}}(x) \text{ contains a long cycle of length } \leq k\right)\\ &\quad\leq \frac{V}{\delta V^{2/3}} \mathbb{P}_{p_c}\left(\mathcal{C}_{\mathbb{T}}(0) \text{ contains a long cycle of length } \leq k\right).\end{aligned}$$

If $\mathcal{C}_{\mathbb{T}}(0)$ contains a long cycle of length at most k, then there exists a vertex $x \in \mathcal{C}_{\mathbb{T}}(0)$ such that the following events occur disjointly: $\{0 \leftrightarrow x \text{ in } \mathbb{T}_r^d\}$ and $\{x \text{ is in a long cycle of length } \leq k\}$. Therefore we can use the BK inequality to bound

$$\frac{V}{\delta V^{2/3}} \sum_{x \in \mathbb{T}_r^d} \mathbb{P}_{p_c}\left(0 \leftrightarrow x \text{ in } \mathbb{T}_r^d\right) \mathbb{P}_{p_c}\left(x \text{ is in a long cycle of length } \leq k\right).$$

The sum $\sum_{x \in \mathbb{T}_r^d} \mathbb{P}_{p_c}\left(0 \leftrightarrow x \text{ in } \mathbb{T}_r^d\right) = \mathbb{E}_{\mathbb{T},p_c}|\mathcal{C}_{\mathbb{T}}(0)|$ is the expected size of $\mathcal{C}_{\mathbb{T}}(0)$. Using (2.5) and translation invariance, we bound

$$\mathbb{P}_{p_c}\left(\exists x\colon |\mathcal{C}_{\mathbb{T}}(x)| > \delta V^{2/3} \text{ and } \mathcal{C}_{\mathbb{T}}(x) \text{ contains a long cycle of length } \leq k\right) \leq \frac{V}{\delta V^{2/3}} C_3 V^{1/3} \mathbb{P}_{p_c}(\mathcal{E}_k).$$

Lemma 4.1 follows by using the upper bound on the probability of $\mathcal{E}_k$ in (4.2). ☐

The next lemma gives a more intuitive sufficient condition for Theorem 1.2.

Lemma 4.2. *Assume (1.1). The following assumption implies Theorem 1.2:*

$$\limsup_{\varepsilon \downarrow 0} \limsup_{n \to \infty} \mathbb{P}_{p_c}\left(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} \partial Q_n \mid 0 \longleftrightarrow \partial Q_n\right) = 0.$$

Proof. We show that the assumption of Lemma 4.2 implies the assumption of Lemma 4.1 for $k = \varepsilon n^2$. Note that there exists $\gamma = \gamma(\varepsilon)$ such that $\gamma > 0$, $\gamma \downarrow 0$ as $\varepsilon \downarrow 0$ and

$$\limsup_{\varepsilon \downarrow 0} \frac{1}{\gamma(\varepsilon)} \limsup_{n \to \infty} \mathbb{P}_{p_c}\left(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} \partial Q_n \mid 0 \longleftrightarrow \partial Q_n\right) = 0.$$

To show that the assumption of Lemma 4.1 holds, we take $k = \varepsilon n^2$ and $n = \gamma(\varepsilon)^{1/2} V^{1/6}$. The result follows, since $\mathbb{P}_{p_c}(0 \leftrightarrow \partial Q_R) \asymp R^{-2}$ by (2.1). ☐

We are now ready to complete the proof of Theorem 1.2.

Proof of Theorem 1.2. The assumption of Lemma 4.2 is satisfied by Theorem 1.5, which shall be proved independently in Section 6 below. Hence, Lemma 4.2 implies Theorem 1.2. ☐

5 Proof of Theorems 1.3 and 1.4

5.1 Proof of Theorem 1.3

Let $\mathcal{I}$ be the set of $z \in \mathcal{C}_{\mathbb{T}}(0)$ such that $\{0 \leftrightarrow z\} \circ \{z \text{ is in a long cycle}\}$. Theorem 1.3 follows from Lemma 5.1:

Lemma 5.1 (Bound on the number of long cycles)**.**

$$Y_{\mathcal{C}_{\mathbb{T}}(0)} \leq 2d \, |\mathcal{I}|.$$

Moreover, when (1.1) holds, there exists a finite constant C such that

$$\mathbb{E}_{p_c} |\mathcal{I}| \leq C V^{-1/3}.$$

Before we prove Lemma 5.1, we show how to use it to complete the proof of Theorem 1.3:

$$\mathbb{E}_{p_c}\left[Y_\delta\right] = \sum_x \mathbb{E}_{p_c}\left[\frac{1}{\mathcal{C}_{\mathbb{T}}(x)} Y_{\mathcal{C}_{\mathbb{T}}(x)} I(|\mathcal{C}_{\mathbb{T}}(x)| > \delta V^{2/3})\right] \leq (\delta V^{2/3})^{-1} V \mathbb{E}_{p_c}\left[Y_{\mathcal{C}_{\mathbb{T}}(0)}\right] \leq C_1/\delta,$$

as required. In the remaining part of this section, we prove Lemma 5.1.

Proof of Lemma 5.1. Let $\mathcal{E}$ be the set of edges of the torus adjacent to at least one of the vertices from $\mathcal{I}$. We will show that $Y_{\mathcal{C}_{\mathbb{T}}(0)} \leq |\mathcal{E}|$.

Let $G = \mathcal{C}_{\mathbb{T}}(0)$, and let $\widetilde{G} = G \setminus \mathcal{E}$ denote the subgraph of G obtained by removing every edge of G that is in $\mathcal{E}$. Note that every vertex from $\mathcal{I}$ is an isolated vertex in $\widetilde{G}$. We claim that the graph $\widetilde{G}$ does not contain long cycles. Indeed, assume that there is a long cycle π in $\widetilde{G}$. Since $\widetilde{G}$ is a subgraph of G, π is a long cycle in G. In particular, there exists $z \in \pi$ such that 0 is connected to z in G by a path that does not use any edges from π. Therefore, $z \in \mathcal{I}$ and z is not an isolated vertex in $\widetilde{G}$. This is a contradiction.

We have just shown that by removing every edge adjacent to a vertex in $\mathcal{I}$, we obtain a subgraph of $\mathcal{C}_{\mathbb{T}}(0)$ without long cycles. This implies that $Y_{\mathcal{C}_{\mathbb{T}}(0)} \leq |\mathcal{E}| \leq 2d|\mathcal{I}|$.

Further, by the BK inequality, (2.5) and Theorem 1.1,

$$\mathbb{E}_{p_c}|\mathcal{I}| \leq \mathbb{E}_{p_c}|\mathcal{C}_{\mathbb{T}}(0)| \; \mathbb{P}_{p_c}(0 \text{ is in a long cycle}) \leq C_2 V^{-1/3}.$$

□

5.2 Proof of Theorem 1.4(a): existence of long cycles

In this section we prove Theorem 1.4(a). We show that there exists $c > 0$ such that

$$\mathbb{P}_{p_c}\left(\text{there exists a long cycle of length } > cV^{1/3}\right) > c.$$

Take $\varepsilon > 0$. The precise value of ε will be determined later. Define

$$M = |\{x \colon x \text{ is in a long cycle of length } > \varepsilon V^{1/3}\}|.$$

Then, clearly,

$$\mathbb{P}_{\mathbb{T},p_c}\left(\exists \text{ a long cycle of length } > \varepsilon V^{1/3}\right) = \mathbb{P}_{\mathbb{T},p_c}(M \neq 0). \tag{5.1}$$

By the second moment method, we can bound

$$\mathbb{P}_{\mathbb{T},p_c}(M \neq 0) \geq \frac{(\mathbb{E}_{\mathbb{T},p_c} M)^2}{\mathbb{E}_{\mathbb{T},p_c} M^2}.$$

We first show that $\mathbb{E}_{\mathbb{T},p_c}M \geq C_3V^{1/3}$, and then that $\mathbb{E}_{\mathbb{T},p_c}M^2 \leq C_4V^{2/3}$. By translation invariance,

$$\mathbb{E}_{\mathbb{T},p_c}M = V\mathbb{P}_{\mathbb{T},p_c}(0 \text{ in a long cycle of length } > \varepsilon V^{1/3}).$$

Let $\mathcal{E}_k = \{0 \text{ in a long cycle of length } \leq k\}$. We write

$$\mathbb{P}_{\mathbb{T},p_c}(0 \text{ in a long cycle of length } > \varepsilon V^{1/3}) \geq \mathbb{P}_{\mathbb{T},p_c}(0 \text{ in a long cycle}) - \mathbb{P}_{\mathbb{T},p_c}(\mathcal{E}_{\varepsilon V^{1/3}}).$$

It follows from Theorem 1.1 that

$$\mathbb{P}_{\mathbb{T},p_c}(0 \text{ in a long cycle}) > C_5V^{-2/3}.$$

Therefore, it remains to show that for some $\varepsilon > 0$,

$$\mathbb{P}_{\mathbb{T},p_c}(\mathcal{E}_{\varepsilon V^{1/3}}) < \frac{C_5}{2}V^{-2/3}. \tag{5.2}$$

By (4.2) with $k = \lfloor \varepsilon V^{1/3} \rfloor$ and $n = \lfloor \delta V^{1/6} \rfloor$ for some $\delta > 0$,

$$\mathbb{P}_{\mathbb{T},p_c}(\mathcal{E}_{\varepsilon V^{1/3}}) \leq C_6\delta^2V^{-2/3} + \mathbb{P}_{\mathbb{Z},p_c}\left(0 \overset{\leq \varepsilon V^{1/3}}{\longleftrightarrow} \partial Q_{\lfloor \delta V^{1/6} \rfloor}\right)^2.$$

It follows from Theorem 1.5 and (2.1) in Theorem 2.1 that there exists $r_0 = r_0(\varepsilon, \delta)$ such that for all $r \geq r_0$ (i.e., $V \geq r_0^d$),

$$\mathbb{P}_{\mathbb{Z},p_c}\left(0 \overset{\leq \varepsilon V^{1/3}}{\longleftrightarrow} \partial Q_{\delta V^{1/6}}\right) \leq C_7\mathrm{e}^{-C_8\delta/\sqrt{\varepsilon}}\delta^{-2}V^{-2/3}.$$

We choose $\delta = \varepsilon^{1/3}$. Then, by taking ε small enough, we deduce (5.2) for all $r \geq r_0$. Therefore,

$$\mathbb{P}_{\mathbb{T},p_c}(0 \text{ in a long cycle of length } > \varepsilon V^{1/3}) \geq \frac{C_5}{2}V^{-2/3}$$

for small enough ε and for all $r \geq r_0 = r_0(\varepsilon)$. We then choose $C_3 = C_3(\varepsilon)$ so that $\mathbb{P}_{\mathbb{T},p_c}(0 \text{ in a long cycle of length } > \varepsilon V^{1/3}) \geq C_3V^{-2/3}$ for all $r \geq 1$. This implies that $\mathbb{E}_{\mathbb{T},p_c}M \geq C_3V^{1/3}$. It remains to prove that $\mathbb{E}_{\mathbb{T},p_c}M^2 \leq C_4V^{2/3}$. Since

$$\mathbb{E}_{\mathbb{T},p_c}M^2 = \sum_{x,y}\mathbb{P}_{\mathbb{T},p_c}\left(x, y \text{ in long cycles of length } > \varepsilon V^{1/3}\right),$$

it suffices to show that

$$\sum_{x,y}\mathbb{P}_{\mathbb{T},p_c}\left(x, y \text{ in long cycles}\right) \leq C_4V^{2/3}.$$

We split the above sum, depending on whether the events $\{x \text{ in long cycle}\}$ and $\{y \text{ in long cycle}\}$ occur disjointly or not. The contribution where these events do occur disjointly can be bounded, using the BK-inequality and Theorem 1.1, by $C_9V^{2/3}$, so that

$$\mathbb{E}_{\mathbb{T},p_c}M^2 \leq C_9V^{2/3} + \sum_{x,y}\mathbb{P}_{\mathbb{T},p_c}\left(x, y \text{ in overlapping long cycles}\right),$$

where the event $\{x, y \text{ in overlapping long cycles}\}$ indicates that all pairs of long cycles, one of which contains x and the other y, share at least one edge.

Lemma 5.2 (Contribution of overlapping cycles)**.** *Assume (1.1). There exists a finite constant C such that*

$$\sum_{x,y} \mathbb{P}_{\mathbb{T},p_c}\,(x, y \textit{ in overlapping long cycles}) \le CV^{2/3}. \tag{5.3}$$

Proof. When x and y are in overlapping long cycles, there exist u, v such that (a) u and v both are part of the long cycle that contains x as well as the one that contains y, (b) the connections $x \leftrightarrow u, u \leftrightarrow v, x \leftrightarrow v, y \leftrightarrow u, y \leftrightarrow v$ all occur disjointly, and (c) at least one of the connections in (b) has displacement at least $r/6$. Therefore, using symmetry and the BK inequality, we can upper bound the sum in (5.3) by

$$\begin{aligned} &4 \sum_{x,y,u,v} \mathbb{P}_{\mathbb{T},p_c}(x \leftrightarrow u \text{ with displacement } \ge r/6)\tau_{\mathbb{T},p_c}(x, v)\tau_{\mathbb{T},p_c}(u, v)\tau_{\mathbb{T},p_c}(y, u)\tau_{\mathbb{T},p_c}(y, v) \\ &\qquad + \sum_{x,y,u,v} \tau_{\mathbb{T},p_c}(x, u)\tau_{\mathbb{T},p_c}(x, v)\mathbb{P}_{\mathbb{T},p_c}(u \leftrightarrow v \text{ with displacement } \ge r/6)\tau_{\mathbb{T},p_c}(y, u)\tau_{\mathbb{T},p_c}(y, v). \end{aligned} \tag{5.4}$$

By Theorem 1.7, we can bound $\mathbb{P}_{\mathbb{T},p_c}(x \leftrightarrow u \text{ with displacement } \ge r/6) \le C_{10}V^{-2/3}$. Let

$$\nabla_{\mathbb{T},p} = \sup_{x,y} \sum_{u,v} \tau_{\mathbb{T},p}(x, u)\tau_{\mathbb{T},p}(u, v)\tau_{\mathbb{T},p}(v, y). \tag{5.5}$$

It follows from [6, Theorem 1.6(iii)] and [14, (1.6)] that $\nabla_{\mathbb{T},p_c} < \infty$. Therefore, we can bound the first sum in (5.4) by

$$C_{10}V^{-2/3} \sum_{x,y,u,v} \tau_{\mathbb{T},p_c}(x, v)\tau_{\mathbb{T},p_c}(u, v)\tau_{\mathbb{T},p_c}(y, u)\tau_{\mathbb{T},p_c}(y, v) \le C_{10}V^{-2/3}V\nabla_{\mathbb{T},p_c}\mathbb{E}_{\mathbb{T},p_c}|\mathcal{C}_{\mathbb{T}}(0)| \overset{(2.5)}{\le} C_{11}V^{2/3}, \tag{5.6}$$

and the second sum in (5.4) by

$$\begin{aligned} &C_{10}V^{-2/3} \sum_{x,y,u,v} \tau_{\mathbb{T},p_c}(x, u)\tau_{\mathbb{T},p_c}(x, v)\tau_{\mathbb{T},p_c}(y, u)\tau_{\mathbb{T},p_c}(y, v) \\ &\quad \le C_{10}V^{-2/3}V \sup_{v} \sum_{x,y,u,v} \tau_{\mathbb{T},p_c}(0, u)\tau_{\mathbb{T},p_c}(u, y)\tau_{\mathbb{T},p_c}(y, v) \sum_{v} \tau_{\mathbb{T},p_c}(v) \\ &\quad = C_{10}V^{-2/3}V\nabla_{\mathbb{T},p_c}\mathbb{E}_{\mathbb{T},p_c}|\mathcal{C}(0)| \overset{(2.5)}{\le} C_{11}V^{2/3}. \end{aligned}$$

These estimates complete the proof. □

We have just shown that $\mathbb{E}_{\mathbb{T},p_c}M^2 \le C_4V^{2/3}$. By the second moment method, we obtain

$$\mathbb{P}_{\mathbb{T},p_c}\left(\exists \text{ a long cycle of length } > \varepsilon V^{1/3}\right) = \mathbb{P}_{\mathbb{T},p_c}(M \neq 0) \ge \frac{(\mathbb{E}_{\mathbb{T},p_c}M)^2}{\mathbb{E}_{\mathbb{T},p_c}M^2} \ge \frac{C_3^2}{C_4} > 0.$$

This completes the proof of the fact that a long cycle of length $> \varepsilon V^{1/3}$ exists with positive probability. □

5.3 Proof of Theorem 1.4(b): non-existence of long cycles

In this section we prove that, for any positive δ, with positive probability uniformly in r, the clusters of size $> \delta V^{2/3}$ do not contain any long cycles. In other words, recalling the definition of Y_δ in Section 5.1, we will prove the following proposition:

Proposition 5.1 (Non-existence of long cycles)**.** *For any positive δ there exists $c > 0$ such that, for all $r \geq 1$,*

$$\mathbb{P}_{\mathbb{T},p_c}(Y_\delta = 0) > c.$$

Proof. For $x \in \mathbb{T}_r^d$, run the following exploration of the edges of $\mathcal{C}_{\mathbb{T}}(x)$ started from x. Enumerate the edges of $\mathbb{T}_r^d$. (In the algorithm we describe now, if there are several edges to choose from, we always pick the edge with the smallest number.) The first stage of the algorithm is the standard depth-first exploration. At this stage, after n steps, the algorithm produces

- the set of explored vertices X_n (which will be a subset of the vertices of $\mathcal{C}_{\mathbb{T}}(x)$),
- the set of explored edges E_n (these will be the explored edges, the occupancy of which we will check),
- the set of open explored edges T_n and the open cluster induced by these edges, also denoted by T_n (which will be part of the depth-first spanning tree of $\mathcal{C}_{\mathbb{T}}(x)$), and
- the set of unexplored edges U_n (the algorithm will not check the occupancy of these edges).

Further, let $W_n = E_n \cup U_n$.

Take $x \in \mathbb{T}_r^d$. Let $X_0 = T_0 = \{x\}$, $W_0 = \varnothing$. Let $n \geq 0$. Assume that X_n, E_n, T_n and U_n are defined. If there is no edge $\{a, b\}$ with $a \in X_n$ and $\{a, b\} \notin W_n$, then we stop the algorithm and write $\mathcal{A}_x = \mathcal{A}_n$ for all $\mathcal{A} \in \{X, E, T, U, W\}$. Otherwise, pick the vertex $a \in X_n$ which is the farthest from x in T_n for which there exists $b \in \mathbb{T}_r^d$ such that $\{a, b\} \notin W_n$. Such a vertex, if it exists, is always *unique*, since we explore depth-first. (We prove this statement in Lemma 5.3(a) at the end of the section.)

Let $e = \{a, b\}$ be the smallest such edge. We distinguish two cases:

1. If $b \notin X_n$, then we define $E_{n+1} = E_n \cup \{e\}$, $U_{n+1} = U_n$, and *check* the occupancy of e.
   (a) If e is open, then we define $X_{n+1} = X_n \cup \{b\}$ and $T_{n+1} = T_n \cup \{e\}$.
   (b) If e is closed, then we define $X_{n+1} = X_n$ and $T_{n+1} = T_n$.
2. If $b \in X_n$ (in this case we call e a *surplus* edge), then we define $X_{n+1} = X_n$, $T_{n+1} = T_n$, $E_{n+1} = E_n$, and $U_{n+1} = U_n \cup \{e\}$, and *do not check* the occupancy of e.

Since the number of edges of $\mathbb{T}_r^d$ is finite, this stage of the algorithm will terminate at some step $N < \infty$. We then write $\mathcal{A}_x = \mathcal{A}_N$ for all $\mathcal{A} \in \{X, E, T, U, W\}$. In particular, X_x is the vertex set of $\mathcal{C}_{\mathbb{T}}(x)$, T_x is the "depth-first" spanning tree of $\mathcal{C}_{\mathbb{T}}(x)$ with root at x, and W_x is the set of edges with at least one end vertex in $\mathcal{C}_{\mathbb{T}}(x)$. The occupancy of edges in E_x is known. In particular, the graph induced by sets of open edges in E_x is T_x. The occupancy of edges in U_x has not been checked. The sets E_x and U_x are disjoint. Also note that, given the set of unexplored edges U_x, the edges in U_x are open independently of each other. An example of an edge $\{a, b\} \in U_x$ is given in Figure 1 below.

We proceed by describing the second stage of the algorithm. The aim of this second stage is to select those surplus edges $\{a, b\}$ that (i) close a long cycle; and (ii) are such that $x \longleftrightarrow b$ is completely disjoint from the long cycle that is created by the addition of the edge $\{a, b\}$; and (iii) there are no long cycles precisely when all these selected edges are closed.

After n steps, the algorithm produces

- the set of open explored edges G_n and the open cluster induced by these edges, also denoted by G_n (which will be a subgraph of $\mathcal{C}_{\mathbb{T}}(x)$ without long cycles),



Figure 1: Example of an edge $\{a,b\} \in U_x$.

- the set of explored edges F_n (which will be a subset of edges of U_x, the occupancy of which we will check), and
- the set of special edges Z_n (the algorithm will not check the occupancy of these edges; each $e \in Z_n$ will have the property that in the graph $G_n \cup \{e\}$, e is in a long cycle π, and there exists a path ρ connecting one of the end-vertices of e to x which is edge disjoint of π).

Note that, according to the first stage of the algorithm, each edge $e \in U_x$ can be written as $\{a,b\}$ such that the unique path from a to x in the spanning-tree T_x passes through b. (We prove this statement in Lemma 5.3(b) at the end of the section.)

Denote by B_x the set of end-vertices with this property, that is, a vertex b is in B_x if and only if there exists a vertex a such that the edge $\{a,b\}$ is in U_x and the unique path from a to x in the spanning tree T_x passes through b.

We enumerate the vertices of B_x subject to the following restriction: a vertex $b \in B_x$ receives a smaller number than $b' \in B_x$ if the unique path from b' to x in the spanning tree T_x passes through b. This ordering of the vertices in B_x can be better understood by introducing an auxiliary abstract tree $\mathbf{T}_x$ rooted at x with the vertex set $\{x\} \cup B_x$ and the following set of oriented (away from the root) edges: For $b, b' \in B_x$, there is an edge from b to b' in $\mathbf{T}_x$, if the unique path from b' to x in the depth-first spanning tree T_x passes through b, and the unique path between b and b' in T_x does not contain any other vertices from B_x. With this definition, we can alternatively say that a vertex $b \in B_x$ has a smaller number than $b' \in B_x$ if there is an oriented path from b to b' in $\mathbf{T}_x$. In other words, we enumerate the vertices of B_x according to their distance to x in the abstract tree $\mathbf{T}_x$. An example of a collection of cycles and the corresponding tree $\mathbf{T}_x$ is given in Figure 2 below.

The second stage of the algorithm goes as follows. Let $G_0 = T_x$, $F_0 = \varnothing$, $Z_0 = \varnothing$ and $B_0 = B_x$. Assume that, for $n \geq 0$, the sets G_n, F_n, Z_n and B_n are defined. If $B_n = \varnothing$, then we stop the algorithm and define $G_x = G_n$, $F_x = F_n$ and $Z_x = Z_n$.

Otherwise, pick a vertex $b \in B_n$ with the biggest number. We distinguish two cases:

1. If there are at least two vertices a and a' such that the edges $\{a,b\}$ and $\{a',b\}$ are in $U_x \setminus (F_n \cup Z_n)$, then define $B_{n+1} = B_n$ and we select the admissible edge with the smallest number. (This is the same numbering of the edges of the torus as in the first stage of the algorithm.)
2. If the vertex a such that the edge $\{a,b\}$ is in $U_x \setminus (F_n \cup Z_n)$ is unique, then we define $B_{n+1} = B_n \setminus \{b\}$ and select this edge.

Assume that the edge $e = \{a,b\}$ is selected.

1. If the graph $G_n \cup \{e\}$ *does not contain* a long cycle, then we define $F_{n+1} = F_n \cup \{e\}$ and $Z_{n+1} = Z_n$ and check the occupancy of e.
   (a) If e is open, then define $G_{n+1} = G_n \cup \{e\}$.



Figure 2: Example of a collection of cycles and the corresponding tree $\mathbf{T}_x$.

(b) If e is closed, then define $G_{n+1} = G_n$.

2. If the graph $G_n \cup \{e\}$ *does contain* a long cycle, then we define $G_{n+1} = G_n$, $F_{n+1} = F_n$ and $Z_{n+1} = Z_n \cup \{e\}$ and *do not check* the occupancy of e. Note that, by the special ordering of the vertices in B_x, every long cycle of the graph $G_n \cup \{e\}$ passes through e and it is *edge disjoint with the unique path from b to x in the tree T_x*.

Since the number of edges of $\mathbb{T}_r^d$ is finite, this stage of the algorithm will terminate at some step $N' < \infty$. We then write $\mathcal{A}_x = \mathcal{A}_{N'}$ for all $\mathcal{A} \in \{G, F, Z\}$. The sets F_x and Z_x are disjoint, and their union is U_x. The occupancy of edges in F_x is known. In particular, the graph induced by set of open edges in $E_x \cup F_x$ is G_x. The occupancy of edges in Z_x has not been checked. In particular, given the set Z_x, the edges in Z_x are open independently of each other. By the definition of Z_x, any edge $e \in Z_x$ is in a long cycle in the graph $G_x \cup \{e\}$, and every long cycle of the graph $G_x \cup \{e\}$ passes through e. Moreover, by the special ordering of the vertices in B_x, any edge $e \in Z_x$ can be written as $\{a, b\}$ so that the unique path from b to x in the spanning tree T_x is edge disjoint from some long cycle of the graph $G_x \cup \{e\}$ (but not necessarily from all long cycles of the graph $G_x \cup \{e\}$).

We run the above defined exploration algorithm for all the open clusters of the torus. We pick a vertex x_1 uniformly on the torus and determine the depth-first spanning tree of the cluster of x_1 with root at x_1, T_{x_1}, the set of explored edges $E_{x_1} \cup F_{x_1}$, and the set of special edges Z_{x_1}. We then pick a vertex x_2 uniformly from the remaining vertices and determine the depth-first spanning tree of $\mathcal{C}_{\mathbb{T}}(x_2)$, T_{x_2}, the set of explored edged $E_{x_2} \cup F_{x_2}$, and the set of special edges Z_{x_2}. We then proceed similarly by selecting $x_3, \ldots, x_M$ and determining T_{x_i}, $E_{x_i} \cup F_{x_i}$ and Z_{x_i}. Here $M = M(\omega)$ is the number of open clusters in the realization ω.

Given the sets of explored edges $E_{x_i} \cup F_{x_i}$, the number of long cycles is defined by the status of the special edges Z_{x_i}. In particular, if all the edges in Z_{x_i} are closed, then $\mathcal{C}_{\mathbb{T}}(x_i)$ does *not* contain long cycles. Note that given the set of explored edges $E_{x_i} \cup F_{x_i}$, the event that all the edges in Z_{x_i} are closed has probability

$$(1 - p_c)^{|Z_{x_i}|}.$$

Also, the size of a cluster is determined by the number of vertices in a spanning tree. Therefore, (remember that M is the number of open clusters in the torus)

$$\mathbb{P}_{p_c}\left(Y_\delta = 0\right) = \mathbb{E}_{p_c}\left[(1-p_c)^{\sum_{i=1}^M |Z_{x_i}| I(|\mathcal{C}(x_i)|>\delta V^{2/3})}\right] \ge (1-p_c)^{\mathbb{E}_{p_c}\left[\sum_{i=1}^M |Z_{x_i}| I(|\mathcal{C}(x_i)|>\delta V^{2/3})\right]}. \tag{5.7}$$

The last step follows from Jensen's inequality.

Let $\mathcal{C}_{(1)}, \ldots, \mathcal{C}_{(M)}$ be all the clusters of the torus sorted from the largest (in the number of vertices) to the smallest with ties broken in an arbitrary way. We will show that

$$\mathbb{E}_{p_c}\left[\sum_{i=1}^M |Z_{x_i}| I(|\mathcal{C}_{\mathbb{T}}(x_i)| > \delta V^{2/3})\right] = \mathbb{E}_{p_c}\left[\sum_{i=1}^M \sum_{x\in\mathcal{C}_{(i)}} \frac{1}{|\mathcal{C}_{(i)}|} |Z_x| I(|\mathcal{C}_{(i)}| > \delta V^{2/3})\right]. \tag{5.8}$$

Fix a percolation realization on the torus. Remember the way we select the vertices $x_1, \ldots, x_M$: select x_1 uniformly on the torus, select x_2 uniformly on $\mathbb{T}_r^d \setminus \mathcal{C}_{\mathbb{T}}(x_1)$, select x_3 uniformly on $\mathbb{T}_r^d \setminus (\mathcal{C}_{\mathbb{T}}(x_1) \cup \mathcal{C}_{\mathbb{T}}(x_2))$, and so on. Given a percolation realization on the torus, we can select the vertices $x_1, \ldots, x_M$ in two steps: first select a permutation σ of $\{1, \ldots, M\}$ (the distribution of σ is irrelevant to us), and then select x_i uniformly from $\mathcal{C}_{(\sigma(i))}$. Note that the sum $\sum_{i=1}^M |Z_{x_i}| I(|\mathcal{C}_{\mathbb{T}}(x_i)| > \delta V^{2/3})$ does not depend on σ, i.e., on the order in which we select clusters, and only depends on which points in clusters we select as x_i's. This implies (5.8).

Note that

$$\mathbb{E}_{p_c}\left[\sum_{i=1}^M \sum_{x\in\mathcal{C}_{(i)}} \frac{1}{|\mathcal{C}_{(i)}|} |Z_x| I(|\mathcal{C}_{(i)}| > \delta V^{2/3})\right] \le \frac{1}{\delta V^{2/3}} \mathbb{E}_{p_c}\left[\sum_{i=1}^M \sum_{x\in\mathcal{C}_{(i)}} |Z_x|\right] = \frac{1}{\delta V^{2/3}} \sum_{x\in\mathbb{T}_r^d} \mathbb{E}_{p_c}[|Z_x|].$$

Therefore, it follows from (5.7) and (5.8) that

$$\mathbb{P}_{p_c}\left(Y_\delta = 0\right) \ge (1-p_c)^{\delta^{-1} V^{1/3} \mathbb{E}_{p_c}[|Z_0|]}.$$

Proposition 5.1 follows once we show that

$$\mathbb{E}_{p_c}[|Z_0|] \le C_{12} V^{-1/3}. \tag{5.9}$$

Recall that $\mathcal{I}$ is the set of $z \in \mathcal{C}_{\mathbb{T}}(0)$ such that $\{0 \leftrightarrow z\} \circ \{z \text{ is in a long cycle}\}$, and let $\mathcal{E}$ be the set of edges with at least one end-vertex in $\mathcal{I}$. By the properties of Z_0, if $e \in Z_0$ is open, then $e \in \mathcal{E}$. Therefore,

$$\mathbb{E}_{p_c}[|Z_0|] = \frac{1}{p_c} \sum_e \mathbb{P}_{p_c}(e \in Z_0, e \text{ is open}) \le \frac{1}{p_c} \mathbb{E}_{p_c}[|\mathcal{E}|] \le \frac{1}{p_c} 2d \mathbb{E}_{p_c}[|\mathcal{I}|].$$

The claim (5.9) now follows from Lemma 5.1. This completes the proof of Proposition 5.1. □

In the remainder of this section, we prove some properties of the exploration algorithm defined in the proof of Proposition 5.1. Remember the notation used in the description of the algorithm.

Lemma 5.3 (Structure depth-first tree). *(a) For $n \ge 0$, let $\widetilde{X}_n$ be the set of vertices $a' \in X_n$ for which there exists $b' \in \mathbb{T}_r^d$ such that the edge $\{a', b'\} \notin W_n$. For each $n \ge 0$, there exists a unique vertex $a \in \widetilde{X}_n$ which is the farthest from x in the tree T_n, and all the other vertices from $\widetilde{X}_n$ belong to the unique path from a to the root x in T_n.*
(b) For all $e \in U_x$, there exist $a, b \in X_x$ such that $e = \{a, b\}$ and the unique path from a to x in the tree T_x passes through b.

Proof. The proof of part (a) is by induction on n. The result is obvious for $n = 0$, since $\widetilde{X}_0 = \{x\}$. Assume that the result holds for all $n' \leq n$. Pick the unique vertex $a \in \widetilde{X}_n$ which is the farthest from x in T_n. Let $\{a, b\} \notin W_n$. (If there are several choices, then we pick the smallest edge according to the numbering of the edges of the torus.) If $\{a, b\}$ satisfies 1(a) of the first stage of the algorithm, then $b \in \widetilde{X}_{n+1}$, and the unique path from b to x in T_{n+1} contains $\widetilde{X}_n$, by the induction assumption. If $\{a, b\}$ satisfies 1(b) or 2 of the first stage of the algorithm, then either $\widetilde{X}_{n+1} = \widetilde{X}_n$ (if there is more than one edge $\{a, b'\} \notin W_n$) or $\widetilde{X}_{n+1} = \widetilde{X}_n \setminus \{a\}$ (if there is the unique edge $\{a, b'\} \notin W_n$). In both cases $\widetilde{X}_{n+1}$ satisfies the statement in part (a) of the lemma, by the induction hypothesis. This completes the proof of (a).

To prove part (b), let $e \in U_x$. There exists $n \geq 0$ such that $U_{n+1} \setminus U_n = \{e\}$. By the definition of the algorithm, there exist a and b such that $e = \{a, b\}$ and a is the farthest vertex in $\widetilde{X}_n$ from x in T_n. Note that the edge e satisfies condition 2 of the first stage of the algorithm. In particular, $b \in X_n$ and $e \notin W_n$. Therefore, $b \in \widetilde{X}_n$. The result in part (b) now follows from part (a) of the lemma. ☐

6 Proof of Theorems 1.5 and 1.6

In this section, we restrict to percolation on $\mathbb{Z}^d$. In particular, all the paths are assumed by default to be in $\mathbb{Z}^d$, and we write here $\{x \leftrightarrow y\}$ for $\{x \leftrightarrow y \text{ in } \mathbb{Z}^d\}$ and $\mathcal{C}(x)$ for $\mathcal{C}_{\mathbb{Z}}(x)$. This section is organized as follows. In Section 6.1, we start with some preparatory lemmas based on the techniques in [19, 20]. We prove Theorem 1.6(a) in Section 6.2, and Theorems 1.6(b) and 1.5 in Section 6.3.

6.1 Preparatory lemmas

The following lemma produces the factor $\sqrt{\varepsilon}$ that is present in Theorems 1.5 and 1.6(b):

Lemma 6.1. *There exists a finite constant C such that for any $\varepsilon > 0$, positive integer n, and $x \in \mathbb{Z}^d$ with $|x| \geq n^2$,*

$$\mathbb{P}_{p_c}\big(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} \partial Q_n, 0 \leftrightarrow x\big) \leq C\sqrt{\varepsilon}\mathbb{P}_{p_c}(0 \leftrightarrow x).$$

Proof of Lemma 6.1. This proof is a slight modification of the proof of [20, Lemma 2.5]. The event $E = \{0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} \partial Q_n\}$ is measurable with respect to $\mathcal{B}_I(\varepsilon n^2) = \{x \in \mathbb{Z}^d : 0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} x\}$. Therefore, [20, Lemma 2.5] implies that for any $x \in \mathbb{Z}^d$ with $|x|$ sufficiently large,

$$\mathbb{P}_{p_c}\left(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} \partial Q_n, 0 \leftrightarrow x\right) \leq C_1\sqrt{\varepsilon n^2 \mathbb{P}_{p_c}(E)}\mathbb{P}_{p_c}(0 \leftrightarrow x).$$

In fact, it follows from the proof of [20, Lemma 2.5] that the above inequality holds for all $x \in \mathbb{Z}^d$ with $|x| \geq n^2$. Finally, remember that $\mathbb{P}_{p_c}(E) \leq C_2 n^{-2}$ by (2.1). This completes the proof. ☐

It follows from Lemma 6.1 and (2.2) that

$$\sum_{x \in Q_{2n^2} \setminus Q_{n^2}} \mathbb{P}_{p_c}\left(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} \partial Q_n, 0 \leftrightarrow x\right) \leq C_3\sqrt{\varepsilon} n^4, \tag{6.1}$$

which shall be used crucially later on.

We next recall some notation from [19]. Recall the definition of a K-regular vertex from [19, Definition 4.1]: For $A \subseteq \mathbb{Z}^d$, let $\mathcal{C}(x; A)$ be the set of vertices y such that $x \leftrightarrow y$ in A. For $x \in \partial Q_n$ and positive integers s and K, we say that x is *s-bad* if $\mathcal{C}(x; Q_n)$ satisfies

$$\mathbb{P}_{p_c}\big(|\mathcal{C}(x) \cap Q_s(x)| < s^4 \log^7 s \mid \mathcal{C}(x; Q_n)\big) \leq 1 - \exp(-\log^2 s).$$

We further say that $x \in \partial Q_n$ is *K-irregular* if there exists $s \geq K$ such that x is s-bad. Otherwise we say that x is *K-regular*.

We say that a pair of vertices (x, y) is (n, K, ε)-admissible if the following conditions hold: (a) $y \in \partial Q_n$ and $x \in Q_{2n^2} \setminus Q_{n^2}$; (b) $0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} y$ in Q_n and $y \leftrightarrow x$; (c) y is K-regular; and (d) the edge $\{y, \widetilde{y}\}$ is pivotal for the event $0 \leftrightarrow x$, where $\widetilde{y}$ is the neighbor of y not in Q_n (if more than one exist, then we choose the first in lexicographical order).

We define $Y(n, K, \varepsilon)$ as the number of (n, K, ε)-admissible pairs. The random variable $Y(n, K, \varepsilon)$ is very similar to $Y(j, K, L)$ defined in the proof of [19, Lemma 5.1].

Remark 2. Note that [19, Lemma 5.1] holds for all $M \geq 1$ and not just for $M \geq L^2/2$ as it is stated. Indeed, [19, Lemma 5.1] follows directly from [19, Lemmas 5.3 and 5.5], both of which hold (and are stated) for all $M \geq 1$.

Lemma 6.2. *There exists a positive constant $C_4 = C_4(K)$ such that for K sufficiently large, any positive integer n and $\varepsilon > 0$,*

$$\mathbb{E}_{p_c} Y(n, K, \varepsilon) \geq C_4(K) n^4 \mathbb{E}_{p_c} |\{y \in \partial Q_n \colon 0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, y \text{ is } K\text{-regular}\}|.$$

Proof. A word for word repetition of the proof of [19, Lemma 5.1] (taking into account Remark 2) gives Lemma 6.2. Indeed, with the notation of [19], the only difference in the proofs arises in the proof (and the statement) of [19, Lemma 5.3], where instead of the event

$$\mathcal{E}_1 = \{0 \leftrightarrow x \text{ in } Q_j, x \text{ is } K-\text{regular and } X_j^{K-reg} = M\},$$

we use the event

$$\widetilde{\mathcal{E}_1} = \{0 \overset{\leq \varepsilon j^2}{\longleftrightarrow} x \text{ in } Q_j, x \text{ is } K-\text{regular and } X_j^{K-reg} = M\}.$$

However, the event $\widetilde{\mathcal{E}_1}$ still can be determined by observing only the edges of $\mathcal{C}(0; Q_j)$. Therefore, the proof of Lemma 5.3 in [19] remains unchanged if we replace the event $\mathcal{E}_1$ with the event $\widetilde{\mathcal{E}_1}$. The proof of Lemma 5.5 in [19] also requires only that the event $\mathcal{E}_1$ must be determined by observing only the edges of $\mathcal{C}(0; Q_j)$, and therefore also holds with $\mathcal{E}_1$ replaced with $\widetilde{\mathcal{E}_1}$. □

Note that for every $x \in Q_{2n^2} \setminus Q_{n^2}$ there exists at most one $y \in \partial Q_n$ such that the pair of vertices (x, y) is (n, K, ε)-admissible. Therefore,

$$\mathbb{E}_{p_c} Y(n, K, \varepsilon) = \sum_{x \in Q_{2n^2} \setminus Q_{n^2}} \mathbb{P}_{p_c}\Big(\exists y \in \partial Q_n \colon (x, y) \text{ is } (n, K, \varepsilon)\text{-admissible}\Big).$$

If (x, y) is (n, K, ε)-admissible, then $\{0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n\}$ and $\{0 \leftrightarrow x\}$ both occur. We use this observation to bound the expected number of (n, K, ε)-admissible pairs from above by

$$\sum_{x \in Q_{2n^2} \setminus Q_{n^2}} \mathbb{P}_{p_c}\left(\exists y \in \partial Q_n \colon 0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, 0 \leftrightarrow x\right) = \sum_{x \in Q_{2n^2} \setminus Q_{n^2}} \mathbb{P}_{p_c}\left(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} \partial Q_n, 0 \leftrightarrow x\right).$$

We can now combine these bounds with the results of (6.1) and Lemma 6.2 to get

$$\mathbb{E}_{p_c}\Big|\{y \in \partial Q_n \colon 0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, y \text{ is } K\text{-regular}\}\Big| \leq \frac{C_3}{C_4(K)}\sqrt{\varepsilon} \tag{6.2}$$

for all large enough K, positive integers n and for all $\varepsilon > 0$ with the constants C_3 from (6.1) (not depending on K, n and ε) and $C_4(K)$ from Lemma 6.2 (not depending on n and ε). We next investigate the contribution from K-irregular y's:

Lemma 6.3. *For all large enough K and for all $\varepsilon > 0$,*

$$\limsup_{n\to\infty} \sum_{y\in\partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\le \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, y \text{ is } K\text{-irregular}\right) \le \frac{1}{2} \limsup_{n\to\infty} \sum_{y\in\partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\le 2\varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n\right).$$

Proof of Lemma 6.3. Recall the definition of an s-locally bad vertex from [19, Definition 4.3]: Let $\mathcal{T}_s^{loc}(x)$ be the event that (a) for all $y \in Q_s(x)$, $|\mathcal{C}(y; Q_{s^{2d}}(x)) \cap Q_s(x)| < s^4 \log^4 s$; and (b) there exist at most $\log^3 s$ disjoint open paths starting in $Q_s(x)$ and ending at $\partial Q_{s^{2d}}(x)$. For $x \in \partial Q_n$ and positive integers s and K, we say that a cluster $\mathcal{C}$ in $Q_{s^{4d^2}}(x) \cap Q_n$ is a "spanning cluster" if (a) $\mathcal{C} \cap Q_n$ intersects both $\partial Q_{s^{4d^2}}(x)$ and $Q_{s^{2d}}(x)$, or (b) $\mathcal{C} = \mathcal{C}(x)$. We say that x is *s-locally bad* if there exist spanning clusters $\mathcal{C}_1, \ldots, \mathcal{C}_m$ in $Q_{s^{4d^2}}(x) \cap Q_n$ such that

$$\mathbb{P}_{p_c}\left(\mathcal{T}_s^{loc}(x) \mid \mathcal{C}_1, \ldots, \mathcal{C}_m\right) \le 1 - \exp(-\log^2 s).$$

Note that the event that x is s-locally bad is determined by the status of the edges in the box $Q_{s^{4d^2}}(x) \cap Q_n$. Moreover, it follows from [19, Claim 4.2] that if x is not K-regular, then there exists $s \ge K$ such that x is s-locally bad. Therefore, we need to bound from above the probabilities

$$\mathbb{P}_{p_c}\left(0 \overset{\le \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, y \text{ is } s\text{-locally bad}\right)$$

for $y \in \partial Q_n$, $s \ge K$ and large enough n.

Since we are only interested in large n, we may assume that $n\varepsilon > 1$. We consider two different cases: $2d(s^{4d^2})^d < n$ and $2d(s^{4d^2})^d \ge n$. We start with the case $2d(s^{4d^2})^d < n$. Note that in this case the ball $Q_{s^{4d^2}}$ contains at most εn^2 edges. We bound the sum

$$\sum_{y\in\partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\le \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, y \text{ is } s\text{-locally bad}\right)$$

from above by

$$\sum_{y\in\partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\le \varepsilon n^2}{\longleftrightarrow} Q_{s^{4d^2}}(y) \text{ in } Q_n, y \text{ is } s\text{-locally bad}\right).$$

Since the events $\{0 \overset{\le \varepsilon n^2}{\longleftrightarrow} Q_{s^{4d^2}}(y) \text{ in } Q_n\}$ and $\{y \text{ is } s\text{-locally bad}\}$ depend on the states of edges in disjoint subsets of $\mathbb{Z}^d$, they are independent. In particular, the above sum equals

$$\sum_{y\in\partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\le \varepsilon n^2}{\longleftrightarrow} Q_{s^{4d^2}}(y) \text{ in } Q_n\right) \mathbb{P}_{p_c}\left(y \text{ is } s\text{-locally bad}\right).$$

By [19, Lemma 1.1] and the FKG inequality (see, e.g., [10, Theorem 2.4]), there exist a positive constant C_5 and a finite constant C_6 such that for all m and $z, z' \in \partial Q_m$,

$$\mathbb{P}_{p_c}(z \leftrightarrow z' \text{ in } Q_m) \ge C_5 \exp\left(-C_6 \log^2 m\right).$$

We apply this result to "extend" the path $0 \overset{\le \varepsilon n^2}{\longleftrightarrow} Q_{s^{4d^2}}(y)$ in Q_n to a path $0 \overset{\le 2\varepsilon n^2}{\longleftrightarrow} y$ in Q_n:

$$\mathbb{P}_{p_c}\left(0 \overset{\le \varepsilon n^2}{\longleftrightarrow} Q_{s^{4d^2}}(y) \text{ in } Q_n\right) \le C_7 \exp\left(C_8 \log^2 s\right) \mathbb{P}_{p_c}\left(0 \overset{\le 2\varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n\right).$$

Here we also use the fact that the number of edges in $Q_{s^{4d^2}}(y)$ is at most εn^2, which implies that if two vertices z and z' in $Q_{s^{4d^2}}(y)$ are connected by an open path in $Q_{s^{4d^2}}(y)$ then the length of this path is at most εn^2.

It follows from [19, Lemma 4.3] that

$$\mathbb{P}_{p_c}\left(y \text{ is } s\text{-locally bad}\right) \leq C_9 \exp\left(-C_{10} \log^4 s\right).$$

We now put these bounds together. Let $\sum'$ be the sum over all s such that $s \geq K$ and $2d(s^{4d^2})^d < n$. We obtain that

$$\sum{}' \sum_{y \in \partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, y \text{ is } s\text{-locally bad}\right)$$

is bounded from above by

$$C_{11} \exp\left(-C_{12} \log^4 K\right) \sum_{y \in \partial Q_n} \mathbb{P}_{p_c}(0 \overset{\leq 2\varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n),$$

where the constants C_{11} and C_{12} do not depend on n, ε or K.

We now consider the case $2d(s^{4d^2})^d \geq n$. Let $s_0 = (n/2d)^{1/(4d^3)}$. For $s \geq s_0$, we simply bound

$$\sum_{s=s_0}^{\infty} \sum_{y \in \partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, y \text{ is } s\text{-locally bad}\right) \leq |\partial Q_n| \sum_{s=s_0}^{\infty} \sup_{y \in \partial Q_n} \mathbb{P}_{p_c}\left(y \text{ is } s\text{-locally bad}\right).$$

We again use [19, Lemma 4.3] to bound the above expression by

$$|\partial Q_n| \sum_{s=s_0}^{\infty} C_9 \exp\left(-C_{10} \log^4 s\right) \leq C_{13} \exp\left(-C_{14} \log^4 n\right),$$

since $s_0 = (n/2d)^{1/(4d^3)}$, and where the constants C_{13} and C_{14} do not depend on n, K, or ε.

We take K so large that $C_{11} \exp\left(-C_{12} \log^4 K\right) < 1/2$. Remember [19, Claim 4.2] which states that if y is K-irregular, then there exists $s \geq K$ such that y is s-locally bad. Therefore, for such choice of K, the sum

$$\sum_{y \in \partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\leq \varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n, y \text{ is } K\text{-irregular}\right)$$

is bounded from above by

$$\frac{1}{2} \sum_{y \in \partial Q_n} \mathbb{P}_{p_c}\left(0 \overset{\leq 2\varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_n\right) + C_{13} \exp\left(-C_{14} \log^4 n\right). \tag{6.3}$$

The result of Lemma 6.3 follows. ☐

With Lemma 6.3 at hand, we are now ready to prove Theorems 1.6 and 1.5 in Sections 6.2 and 6.3.

6.2 Proof of Theorem 1.6(a)

The proof of Theorem 1.6(a) is similar to the proof of Lemma 6.3, but easier. We refer the reader to Section 6.1 for definitions and notation. Remember the definition of a K-regular vertex from Section 6.1. Let X_n^{K-reg} be the number of K-regular vertices on the boundary of Q_n connected to the origin by an open path in Q_n.

Let $Y(n,K,L)$ be the random variable defined in the proof of [19, Theorem 2]: We say that a pair of vertices (x,y) are (n,K,L)-admissible if the following conditions hold: (a) $x \in \partial Q_n$ and $y \in Q_L(x)$; (b) $0 \leftrightarrow x$ in Q_n and $x \leftrightarrow y$; (c) x is K-regular; and (d) the edge $\{x, \widetilde{x}\}$ is pivotal for the event $0 \leftrightarrow y$, where $\widetilde{x}$ is the neighbor of x not in Q_n (if more than one exist, we choose the first in lexicographical order). We define $Y(n,K,L)$ as the number of (n,K,L)-admissible pairs.

It follows from [19, Lemma 5.1] and Remark 2 that there exists $C_{15} = C_{15}(K)$ such that for all large enough K and for all n and L,

$$\mathbb{E}_{p_c} Y(n,K,L) \geq C_{15}(K) L^2 \mathbb{E}_{p_c} X_n^{K-reg}. \tag{6.4}$$

Lemma 6.4. *For all large enough K and for all n,*

$$\mathbb{E}_{p_c} X_n^{K-reg} \geq \frac{1}{3} \sum_{x \in \partial Q_n} \mathbb{P}_{p_c}(0 \leftrightarrow x \textit{ in } Q_n).$$

Proof. The proof of this lemma is very similar to the proof of Lemma 6.3, but simpler. In the same way as in the proof of Lemma 6.3 and with the same choice of K, we bound the sum

$$\sum_{x \in \partial Q_n} \mathbb{P}_{p_c}\left(0 \leftrightarrow x \text{ in } Q_n, x \text{ is not } K\text{-regular}\right)$$

from above by

$$\frac{1}{2} \sum_{x \in \partial Q_n} \mathbb{P}_{p_c}\left(0 \leftrightarrow x \text{ in } Q_n\right) + C_{13} \exp\left(-C_{14} \log^4 K\right).$$

We then use the result of [19, Lemma 3.1]: For all positive integers n,

$$\sum_{x \in \partial Q_n} \mathbb{P}_{p_c}(0 \leftrightarrow x \text{ in } Q_n) \geq 1.$$

We increase K if necessary to fulfill the bound $C_{13} \exp\left(-C_{14} \log^4 K\right) < 1/6$. The result follows. ☐

Theorem 1.6(a) follows in a straightforward way from (6.4), Lemma 6.4 and the fact that

$$\mathbb{E}_{p_c} Y(n,K,n) \leq \sum_{y \in Q_{2n}} \mathbb{P}_{p_c}(0 \leftrightarrow y) \leq C_{16} n^2.$$

The last inequality follows from (2.2). ☐

6.3 Proof of Theorems 1.5 and 1.6(b)

Proof of Theorem 1.6(b). Let

$$F(\varepsilon) = \limsup_{n \to \infty} \sum_{y \in \partial Q_n} \mathbb{P}_{p_c}\left(0 \xleftrightarrow{\leq \varepsilon n^2} y \text{ in } Q_n\right).$$

Theorem 1.6(a) implies that there exists a finite constant C_{17} such that $F(\varepsilon) \leq C_{17}$ for all $\varepsilon > 0$. It follows from (6.2) and Lemma 6.3 that for all $\varepsilon > 0$,

$$F(\varepsilon) \leq \frac{C_3}{C_4}\sqrt{\varepsilon} + \frac{1}{2} F(2\varepsilon).$$

We apply the above inequality k times to get

$$F(\varepsilon) \le C_{18}\sqrt{\varepsilon} + \frac{1}{2^k}F(2^k\varepsilon) \le C_{18}\sqrt{\varepsilon} + \frac{C_{17}}{2^k},$$

with $C_{18} = C_3\sqrt{2}/C_4(\sqrt{2}-1)$ and where we use that $F(2^k\varepsilon) \le C_{17}$ by Theorem 1.6(a). This inequality holds for any fixed k. We complete the proof of Theorem 1.6(b) by taking k such that $2^k\sqrt{\varepsilon} > 1$. ☐

Proof of Theorem 1.5. Take $\varepsilon > 0$ and $n \ge 1$. Let $k = \lfloor K\varepsilon^{1/2}n\rfloor$, for large enough K. The precise choice of K will be made later in the proof. It suffices to prove Theorem 1.5 for small enough ε, therefore, we assume that $K\varepsilon^{1/2} < 1$. (Our choice of K later will be made independently of ε.) Let $M = \lfloor 1/(K\varepsilon^{1/2})\rfloor - 1 > 0$. Note that the event $\{0 \overset{\le\varepsilon n^2}{\longleftrightarrow} \partial Q_n\}$ implies the existence of $x_1, \dots, x_M$ such that for all $i \in \{1, \dots, M\}$, $x_i \in \partial Q_k(x_{i-1})$ (where we assume $x_0 = 0$) and the following connections all occur disjointly:

$$\{x_{i-1} \overset{\le\varepsilon n^2}{\longleftrightarrow} x_i \text{ in } Q_k(x_{i-1})\}, \quad i \in \{1, \dots, M\}, \quad \text{and} \quad \{x_M \leftrightarrow \partial Q_k(x_M)\}.$$

By the BK inequality and translation invariance, we get

$$\mathbb{P}_{p_c}\left(0 \overset{\le\varepsilon n^2}{\longleftrightarrow} \partial Q_n\right) \le \left[\sum_{y\in\partial Q_k} \mathbb{P}_{p_c}\left(0 \overset{\le\varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_k\right)\right]^M \mathbb{P}_{p_c}\left(0 \leftrightarrow \partial Q_k\right).$$

It follows from (2.1) and the definition of k that

$$\mathbb{P}_{p_c}\left(0 \leftrightarrow \partial Q_k\right) \le \frac{C_{19}}{K^2\varepsilon}\mathbb{P}_{p_c}\left(0 \leftrightarrow \partial Q_n\right).$$

Therefore,

$$\limsup_{n\to\infty} \mathbb{P}_{p_c}\left(0 \overset{\le\varepsilon n^2}{\longleftrightarrow} \partial Q_n \mid 0 \leftrightarrow \partial Q_n\right) \le \frac{C_{19}}{K^2\varepsilon}\left[\limsup_{n\to\infty} \sum_{y\in\partial Q_k} \mathbb{P}_{p_c}\left(0 \overset{\le\varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_k\right)\right]^M.$$

By the definition of k and Theorem 1.6(b),

$$\limsup_{n\to\infty} \sum_{y\in\partial Q_k} \mathbb{P}_{p_c}\left(0 \overset{\le\varepsilon n^2}{\longleftrightarrow} y \text{ in } Q_k\right) \le C_{20}/K.$$

The constant C_{20} does not depend on ε. We choose K large enough so that $C_{20}/K < \mathrm{e}^{-1}$. Putting the bounds together, we obtain

$$\limsup_{n\to\infty} \mathbb{P}_{p_c}\left(0 \overset{\le\varepsilon n^2}{\longleftrightarrow} \partial Q_n \mid 0 \leftrightarrow \partial Q_n\right) \le C_{21}\varepsilon^{-1}\mathrm{e}^{-M}.$$

Since $M = \lfloor 1/(K\varepsilon^{1/2})\rfloor - 1$, the result follows. ☐

7 Proof of Theorem 1.7

We prove the first statement of Theorem 1.7 in Lemma 7.1, and the second statement of Theorem 1.7 in Lemma 7.2.

Lemma 7.1. *There exists a finite constant C such that for all $x \in \mathbb{T}_r^d$,*

$$\mathbb{P}_{p_c}(0 \leftrightarrow x \textit{ in } \mathbb{T}_r^d) \le \tau_{\mathbb{Z},p_c}(0,x) + CV^{-2/3}.$$

Proof. Remember the definition of $\Delta_n(x)$ in Section 2.1. Take $x \in \mathbb{T}_r^d$. We write

$$\mathbb{P}_{p_c}(0 \leftrightarrow x \text{ in } \mathbb{T}_r^d) = \mathbb{P}_{p_c}(\Delta_{V^{1/6}}(x)) + \mathbb{P}_{p_c}\left(\{0 \leftrightarrow x \text{ in } \mathbb{T}_r^d\} \setminus \Delta_{V^{1/6}}(x)\right).$$

It follows from Proposition 2.1 and (2.3) that

$$\mathbb{P}_{p_c}(\Delta_{V^{1/6}}(x)) \le \tau_{\mathbb{Z},p_c}(0,x) + C_1 V^{-2/3}.$$

Let $n = \lfloor V^{1/6}/4 \rfloor$. For $x \in \mathbb{T}_r^d$ and $k \ge 1$, we denote by $\mathcal{C}_{\mathbb{T}}(x;k)$ the set of all $y \in \mathbb{T}_r^d$ such that x and y are connected in $\mathbb{T}_r^d$ by an open path with displacement $\le k$. By Proposition 2.1 and Corollary 2.1, the event $\{0 \leftrightarrow x \text{ in } \mathbb{T}_r^d\} \setminus \Delta_{V^{1/6}}(x)$ implies that (a) $\partial \mathcal{C}_{\mathbb{T}}(0;n) \neq \varnothing$, (b) $\partial \mathcal{C}_{\mathbb{T}}(x;n) \neq \varnothing$, and (c) $\mathcal{C}_{\mathbb{T}}(0;n)$ and $\mathcal{C}_{\mathbb{T}}(x;n)$ do not intersect. We refer the reader to Section 2.1 for the basic properties of $\mathcal{C}_{\mathbb{T}}(z;n)$.

Let G be a subgraph of $\mathbb{T}_r^d$. We define $\mathcal{C}_{\mathbb{T}}(0;n;G)$ in the same way as $\mathcal{C}_{\mathbb{T}}(0;n)$, except that we are now only allowed to use edges from G. A coupling similar to the one in Proposition 2.1 gives

$$\sup_G \mathbb{P}_{p_c}\left(\partial \mathcal{C}_{\mathbb{T}}(0;n;G) \neq \varnothing\right) \le \mathbb{P}_{p_c}\left(0 \leftrightarrow \partial Q_n\right).$$

Indeed, the only difference is that now we do not explore edges of $\mathbb{Z}^d$ corresponding to the edges in $\mathbb{T}_r^d \setminus G$.

Let G' be the subgraph of $\mathbb{T}_r^d$ obtained by removing all edges needed to calculate $\mathcal{C}_{\mathbb{T}}(x;n)$. Note that the events (a)-(c) imply that $\partial \mathcal{C}_{\mathbb{T}}(x;n) \neq \varnothing$ and $\partial \mathcal{C}_{\mathbb{T}}(0;n;G') \neq \varnothing$. By conditioning on the set of edges needed to calculate $\mathcal{C}_{\mathbb{T}}(x;n)$ (with $\partial \mathcal{C}_{\mathbb{T}}(x;n) \neq \varnothing$), we obtain that the probability of the intersection of the events (a)-(c) is bounded from above by

$$\sup_G \mathbb{P}_{p_c}\left(\partial \mathcal{C}_{\mathbb{T}}(0;n;G) \neq \varnothing\right) \mathbb{P}_{p_c}\left(\partial \mathcal{C}_{\mathbb{T}}(x;n) \neq \varnothing\right) \le \mathbb{P}_{p_c}\left(0 \leftrightarrow \partial Q_n\right)^2,$$

which is bounded from above by $C_2 n^{-4} \le C_3 V^{-2/3}$ by (2.1). This completes the proof of the lemma. $\square$

Lemma 7.2. *There exists a finite constant C such that for all $x \in \mathbb{T}_r^d$ and for all positive integers n smaller than $r/2$,*

$$\mathbb{P}_{p_c}(0 \leftrightarrow x \textit{ in } \mathbb{T}_r^d \textit{ by a path with displacement } > n) \le Cn^{2-d} + CV^{-2/3}. \tag{7.1}$$

Consequently, there exists a finite constant C such that for all $x \in \mathbb{T}_r^d$,

$$\mathbb{P}_{p_c}(0 \leftrightarrow x \textit{ in } \mathbb{T}_r^d \textit{ by a path with displacement } > r/6) \le CV^{-2/3}. \tag{7.2}$$

Proof. We start with the proof of (7.1). Let $x \in \mathbb{T}_r^d$. Let n be a positive integer smaller than $r/2$. We distinguish two cases: $|x| < 2n/3$ and $|x| \ge 2n/3$. In the first case, we observe that the event $\{0 \leftrightarrow x$ by a path with displacement $> n\}$ implies that there exists a vertex $y \in \partial Q_n$ such that

$$\{0 \leftrightarrow y \text{ in } Q_n\} \circ \{y \leftrightarrow x\}.$$

By the BK inequality, Theorem 1.6(a) and Lemma 7.1,

$$\begin{aligned} &\mathbb{P}_{p_c}(0 \leftrightarrow x \text{ in } \mathbb{T}_r^d \text{ by a path with displacement } > n) \\ &\quad \le \sum_{y \in \partial Q_n} \mathbb{P}_{p_c}(0 \leftrightarrow y \text{ in } Q_n) \mathbb{P}_{p_c}(y \leftrightarrow x) \le C_4 \tau_{\mathbb{Z},p_c}(y,x) + C_4 V^{-2/3}. \end{aligned} \tag{7.3}$$

Since $|x| < 2n/3$, the distance between x and y is at least $n/3$. Therefore, by (1.1), $\tau_{\mathbb{Z},p_c}(y,x) \leq C_5 n^{2-d}$, and (7.1) follows.

In the case $|x| \geq 2n/3$, the event $\{0 \leftrightarrow x \text{ by a path with displacement } > n\}$ implies that there exists a vertex $y \in \partial Q_{\lfloor n/3 \rfloor}$ such that

$$\{0 \leftrightarrow y \text{ in } Q_{\lfloor n/3 \rfloor}\} \circ \{y \leftrightarrow x\}.$$

Similarly to the calculation in the case $|x| < 2n/3$, we obtain

$$\mathbb{P}_{p_c}(0 \leftrightarrow x \text{ in } \mathbb{T}_r^d \text{ by a path with displacement } > n) \leq C_6 \tau_{\mathbb{Z},p_c}(y,x) + C_6 V^{-2/3}.$$

Since, again, y is chosen so that $|x-y| > n/3$, we obtain (7.1). This completes the proof of (7.1) for all $x \in \mathbb{T}_r^d$. To complete the proof of (7.2), we take $n = \lfloor r/6 \rfloor$. Since $r^{2-d} < V^{-2/3} = r^{-2d/3}$ when $d > 6$, (7.2) follows. □

Acknowledgements. We thank Asaf Nachmias for his comments on parts of the manuscript. The work of RvdH and AS was supported in part by the Netherlands Organisation for Scientific Research (NWO). The work of AS was further supported by a grant of the 'Excellence Fund' of Eindhoven University of Technology, as well as by the grant ERC-2009-AdG 245728- RWPERCRI.

References

[1] L. Addario-Berry, N. Broutin, and C. Goldschmidt. (2010) Critical random graphs: limiting constructions and distributional properties. *Electron. J. Probab.*, **15(25)** 741-775.

[2] D. Aldous (1997) Brownian excursions, critical random graphs and the multiplicative coalescent. *Ann. Probab.*, **25** 812-854.

[3] D.J. Barsky and M. Aizenman (1991) Percolation critical exponents under the triangle condition. *Ann. Probab.*, **19** 1520-1536.

[4] B. Bollobás (2001) *Random graphs*, volume **73** of *Cambridge Studies in Advanced Mathematics.* Cambridge University Press, Cambridge, second edition.

[5] C. Borgs, J.T. Chayes, R. van der Hofstad, G. Slade and J. Spencer (2005) Random subgraphs of finite graphs: I. The scaling window under the triangle condition. *Random Structures and Algorithms* **27** 137-184.

[6] C. Borgs, J.T. Chayes, R. van der Hofstad, G. Slade and J. Spencer (2005) Random subgraphs of finite graphs: II. The lace expansion and the triangle condition. *Annals of Probability* **33** 1886-1944.

[7] C. Borgs, J. Chayes, R. van der Hofstad, G. Slade, and J. Spencer (2006) Random subgraphs of finite graphs. III. The phase transition for the n-cube. *Combinatorica*, **26(4)** 395–410.

[8] J.T. Chayes and L. Chayes (1987) On the upper critical dimension of Bernoulli percolation. *Commun. Math. Phys.* **113(1)** 27-48.

[9] P. Erdős and A. Rényi (1960) On the evolution of random graphs. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, **5** 17-61.

[10] G. Grimmett (1999) *Percolation.* Springer-Verlag, Berlin, second edition.

[11] T. Hara (2008) Decay of correlations in nearest-neighbor self-avoiding walk, percolation, lattice trees and animals. *Ann. Probab.* **36(2)** 530-593.

[12] T. Hara, R. van der Hofstad and G. Slade (2003) Critical two-point functions and the lace expansion for spread-out high-dimensional percolation and related models. *Ann. Prob.* **31** 349-408.

[13] T. Hara and G. Slade (1990) Mean-field critical behaviour for percolation in high dimensions. *Commun. Math. Phys.* **128** 333-391.

[14] M. Heydenreich and R. van der Hofstad (2007) Random graph asymptotics on high-dimensional tori. *Comm. Math. Phys.* **270(2)** 335-358.

[15] M. Heydenreich and R. van der Hofstad (2011) Random graph asymptotics on high-dimensional tori II: Volume, diameter and mixing time. *Probab. Theory Related Fields* **149(3-4)** 397-415.

[16] S. Janson, D.E. Knuth, T. Łuczak, and B. Pittel (1993) The birth of the giant component. *Random Structures Algorithms*, **4(3)** 231-358.

[17] S. Janson, T. Łuczak, and A. Rucinski (2000) *Random graphs.* Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley-Interscience, New York.

[18] H. Kesten (1980) The critical probability of bond percolation on the square lattice equals 1/2. *Commun. Math. Phys.* **74(1)** 41-59.

[19] G. Kozma and A. Nachmias (2011) Arm exponents in high-dimensional percolation. *Journal of the American Mathematical Society* **24(2)** 375-409.

[20] G. Kozma and A. Nachmias (2009) The Alexander-Orbach conjecture holds in high dimensions. *Invent. Math* **178(3)** 635-654.

[21] T. Łuczak, B. Pittel, and J. Wierman (1994) The structure of a random graph at the point of the phase transition. *Trans. Amer. Math. Soc.* **341(2)** 721-748.

[22] A. Nachmias and Y. Peres (2008) Critical random graphs: diameter and mixing time. *Ann. Probab.* **36** 1267-1286.

[23] A. Nachmias and Y. Peres (2010) Critical percolation on random regular graphs. *Random Structures Algorithms* **36** 111-148.